

Matrix functions and network analysis

Paola Boito, Federico Poloni

Pisa–Hokkaido–Roma2 Summer School, August–September
2018

Motivation

Functions of matrices: allow to study in the same framework objects such as

- ▶ Matrix power series, e.g., $\exp(A) = I + A + \frac{1}{2}A^2 + \frac{1}{3!}A^3 + \dots$;
- ▶ Maps on eigenvalues, e.g.,

$$A = V\Lambda V^{-1} \mapsto V \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix} V^{-1};$$

- ▶ Matrix rational iterations, e.g., $X_{k+1} = \frac{1}{2}X_k + \frac{1}{2}X_k^{-1}$;
- ▶ Solutions to matrix equations, e.g., $X^2 = A$.

Many objects that appear in applications can be naturally described as functions of matrices — you have probably already encountered $\exp(A)$ and $A^{1/2}$, for instance.

Motivation

On top of this, an overview of two interesting **applications**:

- ▶ Solving certain boundary-value problems / matrix equations appearing in control and queuing theory.
- ▶ Discovering 'important' vertices in a graph (**centrality measures**).

Reference books

- ▶ N. Higham, *Functions of matrices*. SIAM 2008.
- ▶ Golub, Meurant *Matrices, moments, and quadrature*. Princeton 2010 (for the centrality application).
- ▶ Bini, Iannazzo, Meini, *Numerical solution of algebraic Riccati equations*. SIAM 2012. (for the other application).

Polynomials of matrices

Take a scalar polynomial, and evaluate it in a (**square**) matrix, e.g.,

$$p(x) = 1 + 3x - 5x^2 \implies p(A) = I + 3A - 5A^2.$$

Lemma

If $A = S \text{blkdiag}(J_1, J_2, \dots, J_s) S^{-1}$ is a Jordan form, then $p(A) = S \text{blkdiag}(p(J_1), p(J_2), \dots, p(J_s)) S^{-1}$, and

$$p(J_i) = \begin{bmatrix} p(\lambda_i) & p'(\lambda_i) & \dots & \frac{1}{k!} p^{(k)}(\lambda_i) \\ & p(\lambda_i) & \ddots & \vdots \\ & & \ddots & p'(\lambda_i) \\ & & & p(\lambda_i) \end{bmatrix}.$$

Proof Taylor expansion of p at λ_i and powers of shift matrix.

Functions of matrices [Higham book, '08]

We can extend the same definition to arbitrary scalar functions:

Definition

If $A = S \text{blkdiag}(J_1, J_2, \dots, J_s) S^{-1}$ is a Jordan form, then $f(A) = S \text{blkdiag}(f(J_1), f(J_2), \dots, f(J_s)) S^{-1}$, where

$$f(J_i) = \begin{bmatrix} f(\lambda_i) & f'(\lambda_i) & \dots & \frac{1}{k!} f^{(k)}(\lambda_i) \\ & f(\lambda_i) & \ddots & \vdots \\ & & \ddots & f'(\lambda_i) \\ & & & f(\lambda_i) \end{bmatrix}.$$

Given $f : U \subseteq \mathbb{C} \rightarrow \mathbb{C}$, we say that f is **defined on A** if f is defined and differentiable at least $n_i - 1$ times on each eigenvalue λ_i of A . ($n_i = \text{max. size of a Jordan block with eigenvalue } \lambda_i$.)

Reasonable doubt: is it independent of the choice of S ?

Alternate definition: via Hermite interpolation

Definition

$f(A) = p(A)$, where p is a polynomial such that $f(\lambda_i) = p(\lambda_i)$, $f'(\lambda_i) = p'(\lambda_i)$, \dots , $f^{n_i-1}(\lambda_i) = p^{n_i-1}(\lambda_i)$ for each i .

We may use this as a definition of $f(A)$:

- ▶ Does not depend on S ;
- ▶ Does not depend on p .

Obvious from the definitions that it coincides with the previous one.

Remark: be careful when you say “all matrix functions are polynomials”, because p depends on A .

Some properties

- ▶ If the eigenvalues of A are $\lambda_1, \dots, \lambda_s$, the eigenvalues of $f(A)$ are $f(\lambda_1), \dots, f(\lambda_s)$. (geometric multiplicities may decrease)
- ▶ $f(A)g(A) = g(A)f(A) = (fg)(A)$ (since they are all polynomials in A).
- ▶ If $f_n \rightarrow f$ together with 'enough derivatives' (for instance because they are analytic and the convergence is uniform), then $f_n(A) \rightarrow f(A)$.
- ▶ **continuity** If $A_n \rightarrow A$, then $f(A_n) \rightarrow f(A)$.
Proof let p_n be the (Hermite) interpolating polynomial on the eigenvalues of A_n . Interpolating polynomials are continuous in the nodes, so $p_n \rightarrow p$ (coefficient by coefficient). Then
$$\|p_n(A_n) - p(A)\| \leq \|p_n(A_n) - p_n(A)\| + \|p_n(A) - p(A)\| \leq \dots$$

Example: square root

$$A = \begin{bmatrix} 4 & 1 & & \\ & 4 & 1 & \\ & & 4 & \\ & & & 0 \end{bmatrix}, \quad f(x) = \sqrt{x}$$

We look for an interpolating polynomial with

$$p(0) = 0, \quad p(4) = 2, \quad p'(4) = f'(4) = \frac{1}{4}, \quad p''(4) = f''(4) = -\frac{1}{32}.$$

i.e.,

$$\begin{bmatrix} 0 & 0 & 0 & 1 \\ 4^3 & 4^2 & 4 & 1 \\ 3 \cdot 4^2 & 2 \cdot 4 & 1 & 0 \\ 6 \cdot 4 & 2 & 0 & 0 \end{bmatrix} \begin{bmatrix} p_3 \\ p_2 \\ p_1 \\ p_0 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \\ \frac{1}{4} \\ -\frac{1}{32} \end{bmatrix},$$

$$p(x) = \frac{3}{256}x^3 - \frac{5}{32}x^2 + \frac{15}{16}x.$$

Example – continues

$$p(A) = \frac{3}{256}A^3 - \frac{5}{32}A^2 + \frac{15}{16}A = \begin{bmatrix} 2 & \frac{1}{4} & -\frac{1}{64} \\ & 2 & \frac{1}{4} \\ & & 2 & \\ & & & 0 \end{bmatrix}.$$

One can check that $f(A)^2 = A$.

Example – square root

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad f(x) = \sqrt{x}$$

does not exist (because $f'(0)$ is not defined).

(Indeed, there is no matrix such that $X^2 = A$.)

Example – matrix exponential

$$A = S \begin{bmatrix} -1 & & & \\ & 0 & & \\ & & 1 & 1 \\ & & & 1 \end{bmatrix} S^{-1}, \quad f(x) = \exp(x).$$

$$\exp(A) = S \begin{bmatrix} e^{-1} & & & \\ & 1 & & \\ & & e & e \\ & & & e \end{bmatrix} S^{-1}$$

Can also be obtained as $I + A + \frac{1}{2}A^2 + \frac{1}{6}A^3 + \dots$
(not so obvious, for Jordan blocks...)

Example – matrix sign

$$A = S \begin{bmatrix} -3 & & & \\ & -2 & & \\ & & 1 & 1 \\ & & & 1 \end{bmatrix} S^{-1}, \quad f(x) = \text{sign}(x) = \begin{cases} 1 & \text{Re } x > 0, \\ -1 & \text{Re } x < 0. \end{cases}$$

$$f(A) = S \begin{bmatrix} -1 & & & \\ & -1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} S^{-1}.$$

Not a multiple of I , in general.

Instead, we can recover stable / unstable invariant subspaces of A as $\ker(f(A) \pm I)$.

If we found a way to compute $f(A)$ without diagonalizing, we could use it to compute eigenvalues via bisection...

Example – complex square root

$$A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad f(x) = \sqrt{x}$$

We can choose branches arbitrarily: let us say $f(i) = \frac{1}{\sqrt{2}}(1 + i)$,
 $f(-i) = \frac{1}{\sqrt{2}}(1 - i)$.

Polynomial: $p(x) = \frac{1}{\sqrt{2}}(1 + x)$.

$$p(A) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}.$$

(This is the so-called principal square root: we have chosen the values of $f(\pm i)$ in the right half-plane — other choices are possible).

(We get a non-real square root of A if we choose non-conjugate values for $f(i)$ and $f(-i)$)

Example – nonprimary square root

With our definition, if we have

$$A = S \begin{bmatrix} 1 & & \\ & 1 & \\ & & 2 \end{bmatrix} S^{-1}, \quad f(x) = \sqrt{x}$$

we cannot get

$$f(A) = S \begin{bmatrix} 1 & & \\ & -1 & \\ & & \sqrt{2} \end{bmatrix} S^{-1} :$$

either $f(1) = 1$, or $f(1) = -1 \dots$

This would also be a solution of $X^2 = A$, though. This is called a **nonprimary square root** of A . We get nonprimary roots/functions if we choose different branches for Jordan blocks with the same eigenvalue.

Not functions of matrices, with our definition. Also, they are not polynomials in A .

Cauchy integrals

If f is analytic on and inside a contour Γ that encloses the eigenvalues of A ,

$$f(A) = \frac{1}{2\pi i} \int_{\Gamma} f(z)(zI - A)^{-1} dz.$$

Generalizes the analogous scalar formula.

Proof If $A = V\Lambda V^{-1} \in \mathbb{C}^{m \times m}$ is diagonalizable, the integral equals

$$V \begin{bmatrix} \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{z-\lambda_1} dz & & \\ & \ddots & \\ & & \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{z-\lambda_m} dz \end{bmatrix} V^{-1} = V \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_m) \end{bmatrix} V^{-1}.$$

By continuity, the equality holds also for non-diagonalizable A .

Methods

Matrix functions arise in several areas: solving ODEs (e.g. $\exp(A)$), matrix analysis (square roots), physics, ...

Main methods to compute them:

- ▶ Factorizations (eigendecompositions, Schur...),
- ▶ Matrix versions of scalar iterations (e.g., Newton on $x^2 = a$),
- ▶ Interpolation / approximation,
- ▶ Complex integrals.

We will study them in this course. But, first, a detour.

Vectorization

Matrix functions are maps $\mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ (or $\mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$). We introduce some terminology / notation to study **linear** maps between these spaces.

Definition

For $A \in \mathbb{C}^{m \times n}$, $v = \text{vec}(A)$ is the vector $v \in \mathbb{C}^{mn}$ obtained by concatenating the columns of A .

$$\text{vec} \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{bmatrix}.$$

Kronecker products

Definition

Given $M = (m_{ij}) \in \mathbb{C}^{m_1 \times m_2}$, $N \in \mathbb{C}^{n_1 \times n_2}$, the **Kronecker product** $M \otimes N \in \mathbb{C}^{m_1 n_1 \times m_2 n_2}$ is the matrix with blocks $m_{ij}N$.

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \otimes \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = \left[\begin{array}{cc|cc} 1 & 2 & 2 & 4 \\ 0 & 1 & 0 & 2 \\ \hline 3 & 6 & 4 & 8 \\ 0 & 3 & 0 & 4 \end{array} \right].$$

Lemma

$$\text{vec}(AXB) = (B^T \otimes A) \text{vec}(X),$$

i.e., $B^T \otimes A$ is the matrix that represents the linear map $X \mapsto AXB$.

Warning: this is B^T , not B^* (no conjugation).

Properties of Kronecker product

1. Linear in both factors:
 $(\lambda L + \mu M) \otimes N = \lambda(L \otimes N) + \mu(M \otimes N).$
2. $M^* \otimes N^* = (M \otimes N)^*.$
3. $LM \otimes NP = (L \otimes N)(M \otimes P)$, if the dimensions are compatible. Follows from $(AB)X(CD) = A(BXC)D.$
4. $(M \otimes N)^{-1} = M^{-1} \otimes N^{-1}.$
5. Q_1, Q_2 unitary $\implies Q_1 \otimes Q_2$ unitary.
6. If $M = V_1 \Lambda_1 V_1^{-1}$, $N = V_2 \Lambda_2 V_2^{-1}$ are eigendecompositions, then $M \otimes N = (V_1 \otimes V_2)(\Lambda_1 \otimes \Lambda_2)(V_1 \otimes V_2)^{-1}$ is an eigendecomposition.
7. Analogously for SVD, Schur factorization, ...
8. The eigenvalues (singular values) of $M \otimes N$ are the pairwise products of the eigenvalues (singular values) of M and N .

Example: Sylvester equations

Given $A, B, C \in \mathbb{C}^{n \times n}$, find $X \in \mathbb{C}^{n \times n}$ that solves the matrix equation $AX - XB = C$.

When does it have a unique solution?

It is a **linear** system in \mathbb{C}^{n^2} .

$$AX - XB = C \iff (I \otimes A - B^T \otimes I) \text{vec}(X) = \text{vec}(C).$$

If $A = Q_A T_A Q_A^*$, $B^T = Q_B T_B Q_B^*$ are Schur decompositions, then

$$I \otimes A - B^T \otimes I = (Q_A \otimes Q_B)(I \otimes T_A - T_B \otimes I)(Q_A \otimes Q_B)^*$$

is a Schur decomposition.

Hence, $\Lambda(I \otimes A - B^T \otimes I) = (\alpha_i - \beta_j : i, j = 1, \dots, n)$, where $\Lambda(A) = (\alpha_1, \dots, \alpha_n)$, $\Lambda(B) = (\beta_1, \dots, \beta_n)$.

Solution of Sylvester equations

We have proved

Lemma

$AX - XB = C$ has a unique solution iff A and B have no common eigenvalues.

Corollary: $AX - XB = C$ is ill-conditioned if A, B have two close eigenvalues. (It's an **iff** when they are normal.)

Numerical solution: can we beat the naive $O(n^6)$ algorithm “form $I \otimes A - B^T \otimes I$ and treat it as a $n^2 \times n^2$ linear system”?

Yes! [Bartels-Stewart algorithm, 1972].

Idea: invert that Schur decomposition.

- ▶ $(Q_A \otimes Q_B)^* \text{vec}(C)$ equals $\text{vec}(Q_B^* C \overline{Q_A})$
 \rightsquigarrow product in $O(n^3)$.
- ▶ $I \otimes T_A - T_B \otimes I$ has $O(n)$ nonzeros per row
 \rightsquigarrow back-substitution in $O(n^3)$.

Extensions

- ▶ $A \in \mathbb{C}^{m \times m}$, $X \in \mathbb{C}^{m \times n}$, $B \in \mathbb{C}^{n \times n}$: everything works without changes.
- ▶ **Stein's equation** $X - AXB = C$: works analogously. Solvable iff $\alpha_i \beta_j \neq 1$ for all i, j .
- ▶ $AXB - CXD = E$ (generalized Sylvester's equation): works analogously, using generalized Schur factorization $\text{schur}(A, C)$ and $\text{schur}(D, B)$.

Lyapunov equations

$$AX + XA^* = C. \quad (*)$$

They are simply Sylvester equations with $B = -A^*$ (and $C = C^*$). They have a few notable properties.

Lemma

Suppose A has all its eigenvalues in the right half-plane $RHP = \{z \in \mathbb{C} : \operatorname{Re}(z) > 0\}$. Then,

1. (*) has a unique solution.
2. $X = \int_0^\infty \exp(-tA)C \exp(-tA^*) dt$.
3. $X \succ 0$ if $C \succ 0$. (positive definite ordering)

Proof

With $X = \int_0^\infty e^{-tA} C e^{-tA^*} dt$, one has

$$\begin{aligned} AX + XA^* &= \int_0^\infty \left(A e^{-tA} C e^{-tA^*} + e^{-tA} C e^{-tA^*} A^* \right) dt \\ &= \left[-e^{-tA} C e^{-tA^*} \right]_0^\infty = 0 - (-C). \end{aligned}$$

The converse holds, too:

Lemma

If (*) holds with $C \succ 0$ and $X \succ 0$, then A has all its eigenvalues in the RHP.

Proof Let $A^* v = \lambda v$; then,

$$v^* C v = v^* (AX + XA^*) v = \bar{\lambda} v^* X v + \lambda v^* X v = 2 \operatorname{Re}(\lambda) v^* X v.$$

Lyapunov's use of these equations

Proving that certain dynamical systems are stable!

Let $y(t) : [0, \infty] \rightarrow \mathbb{C}^n$ be the solution of $\frac{d}{dt}y(t) = -Ay(t)$.

If I can find $X \succ 0$ and $C \succ 0$ such that $A^T X + XA^T = -C$, then

$$\frac{d}{dt} y(t)^* X y(t) = y(t)^* (-A^* X - XA) y(t) = -y(t)^* C y(t) < 0.$$

\implies The 'energy' $y(t)^* X y(t)$ decreases.