

A separation theorem for entire transcendental maps

Anna Miriam Benini^{*}
 CRM Ennio de Giorgi
 Piazza dei Cavalieri 3
 56100 Pisa, Italy
 ambenini@gmail.com

Núria Fagella[†]
 Dept. de Mat. Aplicada i Anàlisi
 Univ. de Barcelona, Gran Via 585
 08007 Barcelona, Spain
 fagella@maia.ub.es

January 21, 2013

Abstract

We study the distribution of periodic points for a wide class of maps, namely entire transcendental functions of finite order and with bounded set of singular values, or compositions thereof. Fix $p \in \mathbb{N}$ and assume that all dynamic rays which are invariant under f^p land. An interior p -periodic point is a fixed point of f^p which is not the landing point of any periodic ray invariant under f^p . Points belonging to attracting, Siegel or Cremer cycles are examples of interior periodic points. For functions as above, we show that rays which are invariant under f^p , together with their landing points, separate the plane into finitely many regions, each containing *exactly* one interior p -periodic point or one parabolic immediate basin invariant under f^p . This result generalizes the Goldberg-Milnor Separation Theorem for polynomials [GM], and has several corollaries. It follows, for example, that two periodic Fatou components can always be separated by a pair of periodic rays landing together; that there cannot be Cremer points on the boundary of Siegel discs; that “hidden components” of a bounded Siegel disc have to be either wandering domains or preperiodic to the Siegel disc itself; or that there are only finitely many non-repelling cycles of any given period, regardless of the number of singular values.

1 Introduction

Given a holomorphic map $f : \mathbb{C} \rightarrow \mathbb{C}$, we are interested in the dynamical system generated by the iterates of f . In this setup, there is a dynamically meaningful partition of the phase space into two completely invariant subsets: the *Fatou set*, where the dynamics are in some sense stable, and the *Julia set*, where they are chaotic. More precisely, the Fatou set is defined as the open set

$$\mathcal{F}(f) := \{z \in \mathbb{C}; \{f^n\} \text{ is normal in a neighborhood of } z\},$$

^{*}Partially supported by the EU network MRTN-CT-2006-035651 CODY

[†]Partially supported by the catalan grant 2009SGR-792, by the spanish grants MTM-2008-01486, MTM2006-05849 Consolider (including a FEDER contribution) and MTM2011-26995-C02-02, and by the european network MRTN-CT-2006-035651-2-CODY.

and the *Julia set* $J(f)$ as its complement. Another dynamically interesting set is the set of escaping points or *escaping set*

$$I(f) := \{z \in \mathbb{C}; f^n(z) \rightarrow \infty, \text{ as } n \rightarrow \infty\}.$$

The relation between them is that, in general, $J(f) = \partial I(f)$, although for some classes of functions $I(f) \subset J(f)$ ([EL]) and hence $J(f) = \overline{I(f)}$.

In this paper we are mainly concerned with *entire transcendental* maps (abbreviated transcendental maps), that is those entire maps for which infinity is an essential singularity. There are several important differences between the dynamics of transcendental maps and that of polynomials, coming from the very different behavior of iterates in a neighborhood of infinity. For example, while the Julia set of polynomials is always a compact set disjoint from $I(f)$, its analogue for transcendental maps is always unbounded and may contain points of $I(f)$.

Another relevant difference concerns the singularities of the inverse function, which always play a crucial role in holomorphic dynamics. For polynomials (or rational maps) all branches of f^{-1} are well defined in a neighborhood of any point, with the exception of *critical values*, that is, $v = f(c)$ where c is a zero of f' (i.e. a *critical point*). If f is entire, one more type of isolated singularity of f^{-1} is allowed, namely the *asymptotic values*, or points $a \in \mathbb{C}$ for which there is a curve $\gamma(t)$ such that $|\gamma(t)| \rightarrow \infty$ as $t \rightarrow \infty$, and $f(\gamma(t)) \rightarrow a$ as $t \rightarrow \infty$. Informally, asymptotic values have some of their preimages at infinity.

A holomorphic function is a covering outside the set $S(f)$ of *singular values* of f , where

$$S(f) := \overline{\{z \in \mathbb{C}; z \text{ is an asymptotic or critical value for } f\}}.$$

Observe that rational maps, and in particular polynomials, have a finite number of singular values while transcendental maps may have infinitely many such singularities. This is in fact part of the reason why for polynomials, every connected component of the Fatou set (also called *Fatou component*) is preperiodic to the basin of an attracting or parabolic periodic point, or to a periodic rotation domain (Siegel disc). Instead, transcendental maps allow for additional types of Fatou components like *wandering components* (that is non-preperiodic) or *Baker domains* (i.e. periodic components for which all their points tend to infinity under iteration). There exist certain natural classes of transcendental maps for which the dynamics is better understood. One such class is the Eremenko-Lyubich class

$$\mathcal{B} = \{f : \mathbb{C} \rightarrow \mathbb{C} \text{ entire transcendental; } S(f) \text{ is bounded}\}.$$

For functions in class \mathcal{B} , the escaping set is a subset of the Julia set (see [EL], Theorem 1) and therefore there are neither Baker domains nor wandering domains in which the iterates tend to infinity. Recently ([Bi]) has constructed an example of function in class \mathcal{B} with wandering domains whose orbits are unbounded but do not tend to infinity; it is still unknown in general whether there can be functions with wandering domains which have bounded orbits. If $S(f)$ is finite, then f is said to be of *finite type* and has no wandering domains of any kind ([EL], [GK]).

Recall that a function is of *finite order* if $\log \log |f(z)|$ is of the order of $\log |z|$ as $|z| \rightarrow \infty$. For example, $f(z) = e^{z^p}$ has order p . In this paper we restrict to the class

$$\widehat{\mathcal{B}} = \{f = g_1 \circ \dots \circ g_k; k \geq 1, g_i \in \mathcal{B} \text{ and } g_i \text{ has finite order}\}.$$

Observe that class $\widehat{\mathcal{B}}$ is contained in \mathcal{B} and is closed under composition, as is \mathcal{B} , while the class of functions of finite order is not.

The reason why we restrict ourselves to class $\widehat{\mathcal{B}}$, is because it is the most general (natural) class for which (*dynamic*) rays or *hairs* have been proven to exist. Informally, a ray is a maximal unbounded injective curve $g : (0, \infty) \rightarrow I(f)$ such that all its iterates are also unbounded curves with the same properties (see Section 2 for a precise definition). If for some $t_* > 0$, $g(t_*)$ is a singular value or the preimage of a singular value, we redefine the ray to be the curve $g : (t_*, \infty) \rightarrow I(f)$ and call it *broken*. We say that a ray *lands* at $z_0 \in \mathbb{C}$ if $g(t) \rightarrow z_0$ as $t \rightarrow 0$. Observe that broken rays do not land but a ray can land at a singular value. With this definition, rays are always pairwise disjoint.

For polynomials, rays foliate the attracting basin of infinity and belong to the Fatou set (see for example [DH], [Mi]). They are in one to one correspondence with straight rays in the complement of \mathbb{D} via Böttcher coordinates, and therefore their dynamics is governed by $\theta \mapsto d\theta$, where θ is the angle that identifies the ray and d is the degree of the polynomial (recall that a polynomial behaves like $z \mapsto z^d$ in a neighborhood of infinity).

For transcendental maps in class \mathcal{B} , the situation is quite different since rays belong to the Julia set. The existence of rays, and their organization with respect to some symbolic dynamics was settled initially in [DT] for functions of finite type (with some extra technical condition). Recently in [R³S], these results have been extended to the class $\widehat{\mathcal{B}}$ (in fact, to a more general but less natural class than $\widehat{\mathcal{B}}$). We refer to Section 2 for background and precise statements. Of special importance for us are the *p*-periodic rays, i.e., rays that are invariant under f^p . As it is the case for periodic points, *p* is called the *period* of the ray. Unless otherwise specified periods are never assumed to be exact. When the period is 1, the ray is called *invariant* or *fixed*. If two *p*-periodic rays land together at a common point, the curve which consists of the two rays and their common landing point is called a *ray pair* (of period *p*). In contrast, we say that a *p*-periodic ray *lands alone* if its landing point is not the landing point of any other ray of the same period. Observe that, although it is expected to be the case, it is not yet known in general whether rays with different (exact) periods can land together at the same point. Therefore the concept of landing alone depends on the period *p*.

Our goal in the present work is to prove some results concerning the relation between periodic rays, their landing points and the distribution of some special periodic points in the complex plane. More precisely, a *p*-periodic point is called an *interior p-periodic point* if there are no *p*-periodic rays landing at it.

Periodic points which are neither repelling nor parabolic (hence (super)attracting, Siegel or Cremer) are necessarily interior periodic points. However, repelling or parabolic *p*-periodic points can be interior *p*-periodic points as well, for example if they are landing points only of rays which are not fixed by f^p , or if they are not landing points of any ray.

We first restrict the attention to fixed points and fixed rays. As opposed to polynomials, functions in class \mathcal{B} in general have infinitely many of each. We assume that all fixed rays land in \mathbb{C} , and in fact this will be a standing assumption for most of the paper. In the polynomial case, it is enough to assume that the Julia set is connected to ensure that all fixed rays land (in fact that all periodic rays land). For transcendental functions, there is no known necessary and sufficient condition to show that fixed rays land. There are however

many cases in which this can be shown, for example if the postsingular set

$$\mathcal{P}(f) := \overline{\bigcup_{s \in \mathcal{S}(f), n \in \mathbb{N}} \{f^n(s)\}}$$

does not intersect the set of fixed rays (see [Fa], [Rel]).

Let Γ be the graph formed by the fixed rays of f together with their endpoints. A connected component of $\mathbb{C} \setminus \Gamma$ is called a *basic region* for f .

Let us observe some facts which are obvious for polynomials but not even true for transcendental maps. A polynomial of degree d always has $d - 1$ fixed rays (fixed points of $\theta \mapsto d\theta$) and d fixed points, counted with multiplicity. This means that there is always at least one interior fixed point, or one double fixed point, even if all fixed rays land alone (this would be the case with only one basic region). If some fixed rays land together, then more fixed points become multiple or otherwise they become interior as they cannot be landing points of any fixed ray. This elementary counting, and the fact that there are only finitely many basic regions, is possible because of the finiteness of the degree. In the transcendental case there are infinitely many fixed points and infinitely many fixed rays, so a priori one might have infinitely many fixed ray pairs, basic regions and interior fixed points. Our first simple, but rather surprising result is the following proposition.

Proposition A (Finiteness of fixed ray pairs). *Let $f \in \widehat{\mathcal{B}}$ such that all fixed rays land. Then all but finitely many fixed rays land alone at repelling fixed points. In particular, there are only finitely many fixed ray pairs and finitely many basic regions.*

Additionally, a global counting of interior fixed points similar to the polynomial case can be done.

Theorem B (Global counting). *Let $f \in \widehat{\mathcal{B}}$ such that all fixed rays land. Let G be a collection of $L \geq 0$ fixed rays such that it contains the set of rays which do not land alone at repelling fixed points. Then there are exactly $L + 1$ fixed points, counted with multiplicity, which are either landing points of some $g \in G$ or interior fixed points.*

An easy but surprising consequence of this theorem (see Corollary 5.3) is that a function in class $\widehat{\mathcal{B}}$ without fixed ray pairs (but which may have infinitely many singular values) can have at most one interior fixed point, in particular at most one non-repelling fixed point.

To state our main theorem we need to recall the local dynamics around a parabolic fixed point (see [Mi]). In a neighborhood of a fixed point z_0 with multiplicity $m + 1 \geq 2$, a holomorphic map f can be written as $f(z) = z + a(z - z_0)^{m+1} + \mathcal{O}(z - z_0)^{m+2}$. In this case, z_0 is in the boundary of m distinct immediate basins of attraction which are fixed by f , formed by points whose orbits converge to z_0 in a given asymptotic direction. Following Goldberg and Milnor [GM], we call each of these basins a *virtual fixed point* of f . More generally, an immediate basin of attraction for a parabolic point which is fixed by f^p is called a *virtual p -periodic point*.

The following is the main theorem in the paper. For completeness, it also contains the statement of Proposition A.

Main Theorem (Separation Theorem). *Let $f \in \widehat{\mathcal{B}}$ and assume that all fixed rays land. Then there are finitely many basic regions, and each basic region contains exactly one interior fixed point or virtual fixed point.*

It follows that interior fixed points can not be multiple. They can still be parabolic but only if the multiplier is different from 1. Otherwise they would have a virtual fixed point attached contradicting the Separation Theorem.

The same Separation Theorem concerning polynomials with a connected Julia set was proven by Goldberg and Milnor in [GM, Theorem 3.3] and we shall refer to it as the Goldberg-Milnor Theorem. In the polynomial case the finiteness of the set of basic regions is immediate, since there are only finitely many fixed rays and fixed points. In their proof, the finiteness of the degree of f is used in an essential way. In a suitable restriction of each basic region, they construct a weakly polynomial-like map and use the Lefschetz fixed point theory to count the number of critical points and fixed points.

The strategy as it is cannot be adapted to the transcendental setting, where the degree is hardly well defined on any restriction, and where the role of critical values can be carried out also by asymptotic values. However, to prove the Main Theorem we do use Goldberg and Milnor's idea of doing a counting in the basic regions, and we manage to do it successfully by counting directly the fixed points instead of going through critical points.

Like in the polynomial case, the statement about periodic points follows from the fixed case and from the fact that the class $\widehat{\mathcal{B}}$ is invariant under composition. Since this is the most general result and the one that is used in the applications, we state it as a theorem although it follows immediately from the Main Theorem.

Theorem C (Separation Theorem for periodic rays). *Let $f \in \widehat{\mathcal{B}}$ such that p -periodic rays land. Then there are finitely many basic regions for f^p , and each basic region contains exactly one interior p -periodic point or virtual p -periodic point.*

The true power of the Goldberg-Milnor Theorem, and therefore of our Separation Theorem, resides in their remarkable number of corollaries. Some of them were not yet known for transcendental functions of any class. We summarize them in the following statement (see Section 6 for the definition of petals).

Corollary D. *If f is a function in $\widehat{\mathcal{B}}$ whose periodic rays land,*

1. *There are only finitely many interior periodic points of any given period. In particular, there are only finitely many non-repelling cycles of any given period.*
2. *Any two periodic Fatou components can be separated by a periodic ray pair.*
3. *There are no Cremer periodic points on the boundary of periodic Fatou components.*
4. *If z_0 is a parabolic point, for each repelling petal there is at least one ray which lands at z_0 through that repelling petal. In particular, any two disjoint attracting petals invariant under f^p are separated by a ray pair of period p .*

5. For any given period p , there are only finitely many (possibly none) periodic points of period p which are landing points of more than one periodic ray. None of them is the landing point of infinitely many rays of the same period.

Note that statement 1 holds regardless of the presence of infinitely many singular values.

From the fact that Fatou components can be separated by periodic ray pairs we obtain the following additional corollary. Given an invariant Siegel disc Δ , we say that U is a *hidden component* of Δ if U is a bounded connected component of $\mathbb{C} \setminus \overline{\Delta}$. Siegel discs with hidden components have never been found but their existence has never been ruled out either, not even for polynomials.

Corollary E (Hidden components of Siegel discs). *If $f \in \widehat{\mathcal{B}}$ and all periodic rays land, then any hidden component of a bounded invariant Siegel disc is either a wandering domain or preperiodic to the Siegel disc itself.*

This article is structured as follows. In Section 2 we state and prove a result concerning the existence of fixed rays for functions in $\widehat{\mathcal{B}}$ and some expansion properties (Lemma 2.3). We call it the Structural Lemma and it is a consequence of results in [R³S]. Hence, we must introduce logarithmic coordinates and the setup of [R³S] to prove it. We also introduce some additional symbolic dynamics to be able to transfer their statements to the original dynamical plane. This setup has interest on its own and we hope that it can be used for other purposes as well. With the Structural Lemma in hand, in Section 3 we present some results about the topology and general distribution of fixed points in the plane, proving Proposition A. In Section 4 we prove some lemmas about homotopies used in the two following sections, which are dedicated to the proof of Theorem B and to the proof of the Main Theorem respectively. The latter uses the setup and notation of the first so it cannot be read independently. Finally, in Section 7 we prove corollaries D and E.

Acknowledgments

We are thankful to Arnaud Chéritat and to Pascale Roesch for sharing with us the preprint of their recent work on Siegel discs for polynomial with two critical values ([CR]), from which comes the original idea that ended up generating this present paper. We thank Xavier Buff for helpful discussions and we are also thankful to IMUB, to the Complex Dynamics group at Universitat de Barcelona, and to the CODY Marie Curie Network thanks to which this project could be carried out. We give very special thanks to the referee for the very thoughtful comments pointing out many details which were not clear enough. Thanks to them the exposition was largely improved. In addition he/she greatly simplified the proof of Proposition 2.5.

2 Tools and preliminary results

In this section, we present the main features of the dynamical plane for $f \in \widehat{\mathcal{B}}$ and state a Structural Lemma (Lemma 2.3) which is needed for the rest of the paper. We prove it using results of [R³S] together with some study of the combinatorics of f . This last part of the

section is necessary for the proof and interesting in itself, but will not be used later during the paper.

2.1 Structure of the dynamical plane and main technical lemma

The following construction is from [EL] (see also [R³S]). Let $f \in \mathcal{B}$, and D be a Jordan domain containing $S(f), 0$ and $f(0)$. For simplicity we assume that the boundary of D is real analytic. Then the preimage \mathcal{T} of $\mathbb{C} \setminus \overline{D}$ under f consists of countably many unbounded connected components $\{T_\alpha, \alpha \in \mathcal{A}\}$ called *tracts* of f , and $f : T_\alpha \rightarrow \mathbb{C} \setminus \overline{D}$ is a universal covering of infinite degree. If f has finite order then the number of tracts is finite.

Observe that since fibers are discrete, tracts accumulate only at infinity. In particular, only finitely many tracts intersect any compact set.

Lemma 2.1. *There is a simple curve $\delta \subset \mathbb{C} \setminus (\overline{D} \cup \overline{\mathcal{T}})$ connecting \overline{D} to infinity.*

Existence of δ seems to be a rather standard fact for functions with finitely many tracts. However we are not aware of any reference covering the case with infinitely many tracts, so we include a proof here.

Proof. Consider the closure $\widehat{\mathcal{T}}$ of the union of all tracts in the Riemann sphere $\widehat{\mathbb{C}}$. This set is connected and full so its complement in $\widehat{\mathbb{C}}$ is simply connected. By Caratheodory's Theorem, to show that any point in $\partial\widehat{\mathcal{T}}$, including infinity, is accessible by curves in $\widehat{\mathbb{C}} \setminus \widehat{\mathcal{T}}$, it is enough to show that $\partial\widehat{\mathcal{T}}$ is locally connected, or equivalently that $\widehat{\mathcal{T}}$ is locally connected. Local connectivity at interior points is immediate. At any point $z \in \partial\widehat{\mathcal{T}}, z \neq \infty$, it follows from the fact that z is on the boundary of some tract and cannot be accumulated by any other point of $\widehat{\mathcal{T}}$, neither from the same tract (the boundary is a smooth curve), nor from other tracts. To show local connectivity at infinity let $r > 0$ and consider the ball B_r of spherical radius r centered at infinity. Since tracts do not accumulate in any compact set, ∂B_r intersects only finitely many tracts, and $B_r \cap \widehat{\mathcal{T}}$ has only finitely many connected components, exactly one of which contains infinity. It follows that the finitely many components of $B_r \cap \widehat{\mathcal{T}}$ not containing infinity can be removed from B_r giving a simply connected open neighborhood of infinity whose intersection with $\widehat{\mathcal{T}}$ is connected. \square

Observe that, in fact, there are many accesses to infinity, providing plenty of freedom in the choice of δ . In particular, given any two tracts, δ can always be chosen so as to separate them. Let δ be any curve as in Lemma 2.1. The preimages $\{f^{-1}\delta\}$ subdivide each tract T_α into *fundamental domains* $F_{\alpha,i}$ with $i \in \mathbb{Z}$ on which $f : F_{\alpha,i} \rightarrow \mathbb{C} \setminus (\delta \cup \overline{D})$ is univalent (see Figure 1).

When it is not relevant which tract the fundamental domains are contained in, or their position inside the tract, we call a fundamental domain simply F . Observe that for each F , because $f|_F$ is univalent, there is a well defined bijective inverse branch $f_F^{-1} : \mathbb{C} \setminus (\delta \cup \overline{D}) \rightarrow F$.

Definition 2.2 (Dynamic ray). A (*dynamic*) *ray* for f is an injective curve $g : (0, \infty) \rightarrow I(f)$ such that:

- $\lim_{t \rightarrow \infty} |f^n(g(t))| = \infty \quad \forall n \geq 0;$




Figure 1: The disc D , the tracts of f and the fundamental domains.

- $\lim_{n \rightarrow \infty} |f^n(g(t))| = \infty$ uniformly in $[t_0, \infty)$ for all $t_0 > 0$;
- $g(0, \infty)$ is maximal with these properties.

If $g(t_*)$ is a singular value or the preimage of a singular value for $t_* > 0$, then we only consider the curve $g : (t_*, \infty) \rightarrow I(f)$ and we call the ray *broken*.

We say that a ray g is *asymptotically contained* in a tract F if $g(t) \in F$ for all t sufficiently large. Recall that a dynamic ray g is *periodic* (or p -periodic) if $f^p(g) \subset g$ for some $p \geq 1$, and *fixed* if $p = 1$. We say that a dynamic ray *lands* at a point $z_0 \in \mathbb{C}$ if $\lim g(t) = z_0$ as $t \rightarrow 0$. Observe that rays are allowed to land at singular values, but that broken rays do not land.

The following lemma summarizes all of the results from this section that we shall use in the rest of the paper. It relies on the results and constructions from [R³S], however it requires independent proofs that we will give in the following subsections. Recall that maps in class $\widehat{\mathcal{B}}$, as defined in the introduction, are also in class \mathcal{B} . For $R > 0$, we denote by D_R the open disc centered at 0 and of radius R , and by C_R the boundary of D_R .

Lemma 2.3 (Structural Lemma). *Let $f \in \widehat{\mathcal{B}}$. Then*

- Tracts and fundamental domains have a cyclic order at infinity and can be labeled according to that order.*
- Let \mathcal{F} be the union of finitely many fundamental domains. Then for any R sufficiently large, the set $f^{-1}(C_R) \cap \mathcal{F}$ is contained in D_R .*
- For each fundamental domain F , there is a unique fixed dynamic ray g_F which is asymptotically contained in F . Conversely, any fixed ray is asymptotically contained in some fundamental domain.*

The three following subsections are dedicated to the proof of the three statements of Lemma 2.3.

2.2 Logarithmic coordinates and lift of f

For an entire transcendental function in class \mathcal{B} we consider logarithmic coordinates following [EL] (see also [R³S]). The logarithm is a well defined multivalued function on $\mathbb{C} \setminus \overline{D}$. Define the set $\tilde{H} := \exp^{-1}(\mathbb{C} \setminus \overline{D})$ and observe that it contains a right half plane. Also $\exp^{-1}(\delta)$ is a countable family of curves $\tilde{\delta}_i$ dividing \tilde{H} into semi-strips \tilde{S}_i ; without loss of generality we label one of them \tilde{S}_0 and find a branch L_0 of the logarithm mapping $\mathbb{C} \setminus (\overline{D} \cup \delta)$ into \tilde{S}_0 . If we define branches L_n of the logarithm as $L_n(w) = L_0(w) + 2\pi in$, we obtain a consecutive labeling of the strips \tilde{S}_n as images of L_n . Call $\tilde{\delta}_n, \tilde{\delta}_{n+1}$ the preimages of δ bounding the semi-strip \tilde{S}_n . Observe that the tracts do not contain the origin, so the branches of the logarithm are defined on the entire tracts (see Figure 2). The set $\tilde{\mathcal{T}} := \exp^{-1}(\mathcal{T})$ then consists of infinitely many connected components $\tilde{T}_{\alpha,i}$ which do not accumulate in any compact set; the set $\tilde{T}_{\alpha,i}$ is defined as $L_i(T_\alpha)$ so an unbounded part of $\tilde{T}_{\alpha,i}$ is always contained in \tilde{S}_i .




Figure 2: Logarithmic coordinates. The lift \tilde{f} is only defined on the tracts. Each of them is mapped conformally onto the set \tilde{H} .

The following properties hold (see [R³S]):

- \tilde{H} is a $2\pi i$ periodic domain containing a right half plane;
- every $\tilde{T}_{\alpha,i}$ is an unbounded Jordan domain with real parts bounded from below, but unbounded from above (i.e., unbounded to the right);
- the $\tilde{T}_{\alpha,i}$ accumulate only at infinity;
- $\tilde{\mathcal{T}}$ is invariant under translation by $2\pi i$.

Lift of f

Because $f \circ \exp$ is a universal covering on each $\tilde{T}_{\alpha,i}$, it is possible to lift f to a continuous function $\tilde{f} : \tilde{\mathcal{T}} \rightarrow \tilde{H}$ which makes the following diagram commute.

$$\begin{array}{ccc} \tilde{\mathcal{T}} & \xrightarrow{\tilde{f}} & \tilde{H} \\ \exp \downarrow & & \downarrow \exp \\ \mathcal{T} & \xrightarrow{f} & \mathbb{C} \setminus \overline{D} \end{array}$$

On each of the infinitely many connected components $\tilde{T}_{\alpha,i}$ of $\tilde{\mathcal{T}}$, the lift $\tilde{f}|_{\tilde{T}_{\alpha,i}}$ is defined up to translation by multiples of $2\pi i$; so $\tilde{f} : \tilde{\mathcal{T}} \rightarrow \tilde{H}$ is defined up to infinitely many constants. Let us fix any choice of \tilde{f} on each of the tracts $\tilde{T}_{\alpha,0}$, and then choose the remaining constants so that \tilde{f} is $2\pi i$ periodic.

The preimages of $\{\tilde{\delta}_j\}$ under \tilde{f} divide each $\tilde{T}_{\alpha,i}$ into *fundamental domains* $\tilde{F}_{\alpha,i,j}$. Choose the labeling of fundamental domains so that $\tilde{f}(\tilde{F}_{\alpha,i,j}) = \tilde{S}_j$ for each α, i . Because we chose the lift \tilde{f} to be periodic, the labeling on the fundamental domains for \tilde{f} induces a well defined labeling for the fundamental domains for f , in such a way that $\exp(\tilde{F}_{\alpha,i,j}) = F_{\alpha,j}$ for each i (see Figure 3).




Figure 3: Labeling of the fundamental domains.

Vertical order

Because the sets $\tilde{T}_{\alpha,i}$ are unbounded to the right, each $\tilde{T}_{\alpha,i}$ divides any right half plane sufficiently far to the right, into at least two connected components, one of which contains points with arbitrarily large imaginary part (that is said to be *above* $\tilde{T}_{\alpha,i}$ or $\succ \tilde{T}_{\alpha,i}$) and one which contains points with arbitrarily small imaginary part (that is said to be *below* $\tilde{T}_{\alpha,i}$ or $\prec \tilde{T}_{\alpha,i}$). This introduces a natural order \prec on the $\tilde{T}_{\alpha,i}$, called *vertical order at infinity*. We may assume that $\tilde{T}_{\beta,i} \prec \tilde{T}_{\alpha,i}$ if and only if $\beta < \alpha$ (for a fixed i). Observe also that $\tilde{T}_{\alpha,i} \prec \tilde{T}_{\beta,j}$

whenever $i < j$; in this case, the order is just induced by branches of the logarithm. The vertical order on the $\tilde{T}_{\alpha,i}$ induces via the exponential map a *cyclic order at infinity* (still denoted by \prec) on the tracts T_α .

Observe that for any given tract T_α , the fundamental domains $F_{\alpha,i}$ contained in T_α also have a well defined cyclic order; this induces a cyclic order on the set of all fundamental domains.

This proves part (a) of the Structural Lemma 2.3.

2.3 Expansion near infinity

This Section is devoted to prove that, when restricted to a finite number of fundamental domains and far enough to the right, \tilde{f} is uniformly expanding (that is, part (b) of Lemma 2.3). Let C_R be the circle of radius R centered at 0.

Let us recall that the density $\rho_{\mathbb{H}}(z)$ of the hyperbolic metric in the right half plane \mathbb{H} is $\rho_{\mathbb{H}}(z) = \frac{1}{\operatorname{Re} z}$. Also, it is a standard estimate that the hyperbolic density on any simply connected hyperbolic domain Ω is comparable with the inverse of the distance to the boundary. More precisely,

$$\frac{1}{2 \operatorname{dist}(z, \partial\Omega)} \leq \rho_\Omega(z) \leq \frac{2}{\operatorname{dist}(z, \partial\Omega)}.$$

As any tract \tilde{T} in logarithmic coordinates has never vertical width larger than 2π , the maximal distance of a point from the boundary is π , hence on the tract we have the inequality

$$\frac{1}{2\pi} \leq \rho_{\tilde{T}}(x). \quad (2.1)$$

Therefore for any curve γ in \tilde{T} , $\ell_{\text{eucl}}(\gamma) \leq 2\pi \ell_{\tilde{T}}(\gamma)$.

The following result follows from [R³S, Corollary 5.8 (a) and (b)]. (Compare also with [R³S, Definition 5.1]).

Proposition 2.4. *If \tilde{f} is a lift of a function $f \in \widehat{\mathcal{B}}$, then the tracts of \tilde{f} satisfy the bounded slope condition, which means that there exists M_1, M_2 such that for any z, w belonging to the same tract,*

$$|\operatorname{Im} z - \operatorname{Im} w| \leq M_1 \max\{\operatorname{Re} z, \operatorname{Re} w, 0\} + M_2. \quad (2.2)$$

We are now ready to prove the following proposition.

Proposition 2.5 (Almost straight). *Let $f \in \widehat{\mathcal{B}}$, and \mathcal{F} be a finite union of fundamental domains for f . Then for any R sufficiently large, $f^{-1}(C_R) \cap \mathcal{F}$ is contained in the bounded connected component of $\mathbb{C} \setminus C_R$.*

Proof. Observe that it is enough to prove the claim for one fundamental domain F . Recall the notation from Section 2.2, and let \tilde{f} be the lift of f defined on the tracts $\tilde{T}_{\alpha,i}$ with the conventions adopted there. Let \tilde{F} be any fundamental domain for \tilde{f} such that $\exp(\tilde{F}) = F$; it is then sufficient to prove that for any R' sufficiently large, and any z with $\operatorname{Re} z = R'$, we have that $\operatorname{Re} u < R'$ for all $u \in \tilde{f}^{-1}(z) \cap \tilde{F}$. The proposition would then hold with $R > e^{R'}$.

Let \tilde{T} be the tract containing \tilde{F} , and let $\tilde{S} = \tilde{f}(\tilde{F})$. Since $\tilde{f} : \tilde{T} \rightarrow \tilde{H}$ is a biholomorphism, there is a well defined inverse $\tilde{f}_{\tilde{T}}^{-1}$, which is an isometry between the hyperbolic metric on \tilde{H} and the hyperbolic metric in \tilde{T} . Its restriction $\tilde{f}_{\tilde{F}}^{-1} : \tilde{S} \rightarrow \tilde{F}$ is hence also an isometry between the hyperbolic metric on \tilde{H} and the hyperbolic metric in \tilde{T} .

Let us first show that for any $z, w \in \tilde{S}$ with $\operatorname{Re} w > \operatorname{Re} z > 0$,

$$|\operatorname{Im} z - \operatorname{Im} w| \leq M_1 \operatorname{Re} w + M_2 + 2\pi, \quad (2.3)$$

where M_1, M_2 are the constants from Proposition 2.4. Indeed, since $\operatorname{Re} w > \operatorname{Re} z$ and tracts are connected sets containing points of arbitrary large real part, there is a point w' in the same tract as z with $\operatorname{Re} w' = \operatorname{Re} w$, for which Equation (2.2) holds. Since $w, w' \in \tilde{S}$ and $\operatorname{Re} w' = \operatorname{Re} w$, we have that $|\operatorname{Im} w - \operatorname{Im} w'| \leq 2\pi$ and (2.3) follows.

Let $\mathbb{H}_L = \{z \in \mathbb{C} \mid \operatorname{Re} z > L\}$ be a right half plane such that $\mathbb{H}_L \subset \tilde{H}$. Then $\rho_{\mathbb{H}_L} \geq \rho_{\tilde{H}}$. Let z_0 be a point in $\tilde{S} \cap \mathbb{H}_L$, and z'_0 be its unique preimage in \tilde{F} . We will show the claim by proving that for any $R' > 0$, and any point $z \in \tilde{H}$ with $\operatorname{Re} z = R' + \operatorname{Re} z_0$, we have $|z - z_0| \geq R'$ while $|\tilde{f}^{-1}(z) \cap \tilde{F} - z'_0| \sim \ln R'$. By direct computation, and using (2.3),

$$\begin{aligned} \operatorname{dist}_{\tilde{H}}(z_0, z) &\leq \int_{\operatorname{Re} z_0}^{\operatorname{Re} z} \frac{1}{t - L} dt + \left| \int_{\operatorname{Im} z_0}^{\operatorname{Im} z} \frac{1}{\operatorname{Re} z - L} dt \right| = \ln \left(\frac{R' + \operatorname{Re} z_0 - L}{\operatorname{Re} z_0 - L} \right) + \frac{|\operatorname{Im} z_0 - \operatorname{Im} z|}{R' + \operatorname{Re} z_0 - L} \leq \\ &\leq \ln R' + k, \end{aligned}$$

where k is a constant. Since \tilde{f} is an isometry for the hyperbolic metric between \tilde{H} and \tilde{T} , we have that

$$\operatorname{dist}_{\tilde{T}}(z'_0, \tilde{f}^{-1}(z) \cap \tilde{F}) \leq \ln R' + k,$$

and by Equation (2.1), $|z'_0, \tilde{f}^{-1}(z) \cap \tilde{F}| \leq 2\pi(\ln R' + k)$. \square

This concludes the proof of part (b) in Lemma 2.3.

2.4 Dynamic rays and their properties

Here we state a result from [R³S] about existence of dynamic rays for \tilde{f} (recall that \tilde{f} is only defined on \tilde{T}). Then, we prove a theorem about existence of dynamic rays for f corresponding to part (c) of the Structural Lemma 2.3.

We define the escaping set

$$I(\tilde{f}) := \{\tilde{z} \in \tilde{T}; \operatorname{Re} \tilde{f}^n(\tilde{z}) \rightarrow \infty \text{ as } n \rightarrow \infty\}.$$

The following definition of ray tail and dynamic ray is from [R³S, Definition 2.2].

Definition 2.6. Let \tilde{f} be the lift of a function f in class $\hat{\mathcal{B}}$. A *ray tail* for \tilde{f} is an injective curve $\tilde{G} : [t_0, \infty) \rightarrow I(\tilde{f})$ such that $\lim_{t \rightarrow \infty} \operatorname{Re} \tilde{f}^n(\tilde{g}(t)) = +\infty$ for any $n > 0$ and such that $\operatorname{Re} \tilde{f}^n(\tilde{g}(t)) \rightarrow +\infty$ uniformly in t as $n \rightarrow \infty$.

Likewise ray tails can be defined for f . Observe that with this definition a non-broken dynamic ray for f becomes a maximal injective curve $g(t) : (0, \infty) \rightarrow I(f)$ such that $g|_{[t, \infty)}$

is a ray tail for every $t > 0$. Observe also that if a point belongs to a ray tail g , by definition it also belongs to a dynamic ray containing g .

Let us define

$$J(\tilde{f}) := \{\tilde{z} \in \tilde{\mathcal{T}} \mid \tilde{f}^n(\tilde{z}) \text{ is defined for all } n\}.$$

Observe that $J(\tilde{f})$ includes orbits which escape to infinity but also any periodic orbit whose projection does not intersect D .

For points $\tilde{z} \in J(\tilde{f})$, we can naturally introduce symbolic dynamics using the labeling of the tracts (see [DT], [R³S],[Ba]): if for $n \geq 0$ we have $\tilde{f}^n(\tilde{z}) \in \tilde{T}_{\alpha_n, i_n}$, we say that \tilde{z} has *address*

$$\tilde{s} := \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \cdots \binom{\alpha_n}{i_n} \cdots$$

and introduce the left-sided shift map σ as

$$\sigma : \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \cdots \mapsto \binom{\alpha_1}{i_1} \binom{\alpha_2}{i_2} \cdots$$

A ray tail $\tilde{G}_{\tilde{s}}$ for \tilde{f} is called a ray tail *of address* \tilde{s} if it consists of points with address \tilde{s} , and analogously for f . We define the set

$$J_{\tilde{s}}(\tilde{f}) := \{\tilde{z} \in J(\tilde{f}) \mid \tilde{z} \text{ has address } \tilde{s}\}$$

From the definition of $J_{\tilde{s}}$, it follows that $\tilde{f}(J_{\tilde{s}}) \subset J_{\sigma\tilde{s}}$; in particular, if an address \tilde{s} is periodic of period m , we have that $\tilde{f}^m(J_{\tilde{s}}) \subset J_{\tilde{s}}$.

The next theorem is the restatement of one of the main results from [R³S].

Theorem 2.7 (Ray tails for \tilde{f}). *Let \tilde{f} be the lift of a function $f \in \hat{\mathcal{B}}$. Then for any address \tilde{s} for which $J_{\tilde{s}}(\tilde{f}) \cap I(\tilde{f}) \neq \emptyset$, the set $J_{\tilde{s}}(\tilde{f})$ contains a unique maximal unbounded arc consisting of escaping points. Any unbounded arc $\tilde{G}_{\tilde{s}} : [0, \infty) \rightarrow J_{\tilde{s}}(\tilde{f})$ is a ray tail of address \tilde{s} for \tilde{f} .*

Theorem 2.7 is not stated in this terms in [R³S], so we sketch a proof using results and terminology from the source.

Proof. From Theorem 5.6 and Lemma 5.7 in [R³S], \tilde{f} satisfies a uniform linear head start condition (uniform head start and uniform bounded wiggling are equivalent by Proposition 5.4). By Theorem 3.3 and Proposition 4.4 (b) in [R³S] there exists a constant $K(\tilde{f})$ such that if $K > K(\tilde{f})$ and the set

$$J_{\tilde{s}}^K(\tilde{f}) := \{\tilde{z} \in J_{\tilde{s}}(\tilde{f}) \mid \operatorname{Re} \tilde{f}^n(\tilde{z}) \geq K, \text{ for all } n \in \mathbb{N}\} \quad (2.4)$$

is non empty, then $J_{\tilde{s}}(\tilde{f})$ has a unique unbounded component, which is a closed arc tending to infinity on one side. From their Corollary 4.4, this arc consists of escaping points and $J_{\tilde{s}}^K(\tilde{f}) \neq \emptyset$. So any unbounded arc $\tilde{G}_{\tilde{s}} : [0, \infty) \rightarrow J_{\tilde{s}}(\tilde{f})$ satisfies the hypothesis of being a ray tail, and consists of points of address \tilde{s} . \square

We now establish a correspondence between existence of ray tails for \tilde{f} and existence of ray tails for f . This discussion can be seen as a natural follow up of the results in [R³S] but it is not contained there.

Given a point z in the f -plane, whose iterates always belong to \mathcal{T} , and such that $f^n(z) \in F_{\alpha_n, i_n}$, we define its *address* s as

$$s = \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \dots .$$

The left-sided shift map σ is well defined, and a point with address s is mapped to a point with address σs .

Lemma 2.8 (Correspondence of addresses). *Let $f \in \widehat{\mathcal{B}}$, \tilde{f} be a periodic lift of f as described in Section 2.2. A point $\tilde{z} \in J(\tilde{f})$ has address $\tilde{s} = \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \dots$ if and only if $z = \exp(\tilde{z})$ has address $s = \binom{\alpha_0}{i_1} \binom{\alpha_1}{i_2} \dots$*

Proof. If $\tilde{z} \in \tilde{T}_{\alpha_n, i_n}$ and $\tilde{f}(\tilde{z}) \in \tilde{T}_{\alpha_{n+1}, i_{n+1}} \subset \tilde{S}_{i_{n+1}}$, then $\tilde{z} \in \tilde{F}_{\alpha_n, i_n, i_{n+1}}$ by the choice of lift \tilde{f} . Hence $z = \exp(\tilde{z}) \in F_{\alpha_n, i_{n+1}}$ by the choice of labeling for the fundamental domains. For the converse, observe that $\exp^{-1}(F_{\alpha_n, i_{n+1}}) = \{\tilde{F}_{\alpha_n, m, i_{n+1}}\}_{m \in \mathbb{Z}}$ and $\tilde{F}_{\alpha_n, m, i_{n+1}} \subset \tilde{T}_{\alpha_n, m}$. \square

Proposition 2.9 (Rays for f). *Let $f \in \widehat{\mathcal{B}}$, \tilde{f} be a periodic lift of f as described in Section 2.2. If $s = \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \dots$ is an address for f , there is a dynamic ray g_s which contains the image under the exponential map of any of the ray tails $g_{\tilde{s}_m}$ (whenever they exist), where $\tilde{s}_m = \binom{\alpha_0}{m} \binom{\alpha_1}{i_0} \dots$, and $m \in \mathbb{Z}$.*

Proof. The image of any ray tail $G_{\tilde{s}_m}$ under the exponential map is a ray tail G_s for f , consisting of points of address s by Lemma 2.8. Then any point in G_s is contained in the dynamic ray g_s of address s by definition hence $g_s \neq \emptyset$. \square

The next proposition ensures that rays for f exist for any address s which contain only finitely many symbols, by showing that the set $J_s^K(\tilde{f})$ defined in Equation 2.4 is non empty. The proof uses ideas from [DT], results from [R³S], and Lemma 2.3 (b).

Proposition 2.10. *If $f \in \widehat{\mathcal{B}}$ and s is an address which contains only finitely many different symbols then there exists a ray g_s with address s for f .*

Proof. Let $s = \binom{\alpha_0}{i_0} \binom{\alpha_1}{i_1} \dots$, and let $J_s^K(f)$ be the set of points of address s such that $|f^n(z)| \geq e^K$ for all $n \in \mathbb{N}$. By definition of \tilde{f} and by Lemma 2.8, $J_s^K(f) = \exp J_{\tilde{s}}^K(\tilde{f})$ (as defined in Equation 2.4) for any $\tilde{s} = \binom{\alpha_0}{m} \binom{\alpha_1}{i_0} \dots$, $m \in \mathbb{N}$. By Theorem 3.3 and Proposition 4.4 (b) in [R³S] (see also the proof of Theorem 2.7 in this paper) there exists $K(\tilde{f})$ such that if $K > K(\tilde{f})$ and $J_{\tilde{s}}^K(\tilde{f})$ is not empty then a ray tail of address \tilde{s} exists. We start by showing that for some $K > K(\tilde{f})$ the set $J_s^K(f)$ is not empty provided s contains only finitely many symbols. By definition a point z has address s if and only if $f^n(z) \in F_{\alpha_n, i_n}$. To simplify notation let us call $F_n := F_{\alpha_n, i_n}$. If s contains only finitely many different symbols, all the F_n belong to some finite collection \mathcal{F} of fundamental domains. Let $R > e^{K(\tilde{f})}$ be such that

Proposition 2.5 holds for the family \mathcal{F} . Let C_R be the circle of radius R , D_R be the open disc of radius R , both centered at 0. Note that $F_n \cap C_R \neq \emptyset$ for all $F_n \in \mathcal{F}$. For each $F_n \in \mathcal{F}$ define ψ_n as the unique inverse branch of f mapping $\mathbb{C} \setminus (\overline{D} \cup \delta) \rightarrow F_n$. By definition each ψ_n is univalent. All ψ_n also have the following property. Let Y be any unbounded connected set; since ψ_n is continuous and it is the inverse of an entire function, its image $\psi_n(Y)$ is also an unbounded connected set. Then, if $Y \cap C_R \neq \emptyset$, we also have that $\psi_n(Y) \cap C_R \neq \emptyset$ because by Proposition 2.5 $\psi_n(Y) \cap D_R \neq \emptyset$. We refer to this property as ψ_n *preserves unboundedness and intersection with C_R* .

For each $N \in \mathbb{N}$ define the set

$$X_N = \{z \in F_0; f^j(z) \in F_j \text{ and } |f^j(z)| \geq R \forall j = 0 \dots N\}.$$

By construction, $\overline{X}_{N+1} \subset X_N$, where the closure is taken in \mathbb{C} . We now show that $\bigcap_N X_N \neq \emptyset$ by additionally showing that $X_N \cap C_R \neq \emptyset$ for all N . Observe that for each N the set X_N is obtained by intersecting F_N with $\mathbb{C} \setminus D_R$, then applying ψ_{N-1} , then intersecting again with $\mathbb{C} \setminus D_R$, applying ψ_{N-2} and so on. Fix N . Since F_N is an unbounded connected set and C_R is compact, $F_N \setminus D_R$ contains at least one unbounded connected component which intersects C_R by definition. So since ψ_{N-1} preserves unboundedness and intersection with C_R , the set $\psi_{N-1}(F_N \setminus D_R)$ also contains at least one unbounded connected component intersecting C_R . By induction, this property is preserved at each step and X_N also contains an unbounded connected component intersecting C_R . Observe that $X_{N+1} \cap C_R$ is compactly contained in $X_N \cap C_R$, so that $\bigcap_N X_N \neq \emptyset$.

Now let $z \in \bigcap_N X_N$. Since $R > e^K$, $\bigcap_N X_N \subset J_s^K(f)$. By definition z has address s , so by Lemma 2.8 (observe that having an address implies that $f^n(z) \in \mathcal{T}$ for all $n \in \mathbb{N}$) there exists a point $\tilde{z} \in \exp^{-1} z$ with address $\tilde{s} = \binom{\alpha_0}{0} \binom{\alpha_1}{i_0} \dots$ such that $\operatorname{Re} \tilde{f}^n(z) \geq K \geq K(\tilde{f})$ for all n . So $J_{\tilde{s}}^K(\tilde{f}) \neq \emptyset$ and there exists a ray tail for \tilde{f} with address \tilde{s} . By Proposition 2.9, there exists a ray for f with address s . \square

Finally we show that each fundamental domain F contains asymptotically exactly one fixed ray.

Proposition 2.11 (Fixed rays in fundamental domains). *For each fundamental domain F , there is a unique fixed dynamic ray g_F such that $g_F(t) \in F$ for all sufficiently large t .*

Proposition 2.11 is a corollary of Theorem 2.7, Proposition 2.10, and the following lemma.

Lemma 2.12. *Any dynamic ray is asymptotically contained in some fundamental domain.*

Proof. Suppose that a ray g is not asymptotically contained in any fundamental domain. Then for some sequence $t_n \rightarrow \infty$, $g(t_n)$ belongs to preimages of δ . Hence $f(g)$ intersects δ infinitely many times and $f^2(g)$ intersects D infinitely many times as $t \rightarrow \infty$, contradicting the definition of dynamic ray. \square

This concludes the proof of the Structural Lemma 2.3, part (c).

3 Location of interior fixed points. Proof of Proposition A

Let us now fix a function $f \in \widehat{\mathcal{B}}$, and restrict our discussion to fixed points and fixed rays. From now on, we assume that all fixed rays land. Let D , δ , etc. be as in Section 2.1 (see Figure 1).

We first prove a proposition about landing of fixed rays for fundamental domains which do not intersect D , and then collect some remarks about the structure of the plane in logarithmic coordinates and the distribution of the interior fixed points with respect to the partition of the plane induced by the fixed rays.

As tracts and fundamental domains accumulate only at infinity, only finitely many fundamental domains can intersect D . On those not intersecting D , the dynamics is easy to study. The following proposition is central to prove Proposition A.

Proposition 3.1 (Forced landing). *If a fundamental domain F does not intersect D , it contains a unique fixed point w , which is repelling, and the fixed ray g_F given by part (c) in the Structural Lemma 2.3 lands at w .*

Proof. Consider a circle C_R of radius R such that part (b) of the Structural Lemma 2.3 holds, and let D_R be the bounded connected component of $\mathbb{C} \setminus C_R$. As $f : F \rightarrow \mathbb{C} \setminus (\overline{D} \cup \delta)$ is univalent, there is a branch ψ of f^{-1} mapping $D_R \setminus (\overline{D} \cup \delta)$ to a proper subset of $F \cap (D_R \setminus (\overline{D} \cup \delta))$ by the Structural Lemma part (c) (see Figure 4). By the Schwarz Lemma, ψ is strictly contracting and has a unique attracting fixed point $w \subset F$. Hence, under iterates of ψ every point tends to a unique fixed point w which is repelling for f ; in particular, any invariant curve converges to w . \square




Figure 4: If F does not intersect D , the fixed ray g_F lands at the unique fixed point in F which must be repelling. However fixed rays from other fundamental domains which do intersect D , can also land at the same fixed point. There are only a finite number of those freelanders though. Light shaded we see the image of the dark shaded domain.

Remark 3.2. Observe that the ray g_F does not necessarily land alone; other fixed rays coming from fundamental domains which do intersect the disc might land together with g_F (see Figure 4).

Recall that a fixed point is an *interior fixed point* if there are no fixed rays landing at it. Observe that interior fixed points cannot be in $\mathbb{C} \setminus (\overline{D} \cup \mathcal{T})$, because f maps $\mathbb{C} \setminus (\overline{D} \cup \mathcal{T})$ to \overline{D} . Also, they cannot be in a fundamental domain not intersecting the disc by Proposition 3.1. So interior fixed points are contained either in $\overline{D} \setminus \mathcal{T}$ or in the finitely many fundamental domains which intersect D (see Figure 5).




Figure 5: Interior fixed points can only lie in the finite number of fundamental domains which intersect D , or in $\overline{D} \setminus \mathcal{T}$.

Recall that Γ is the graph formed by fixed rays together with their endpoints, and that the connected components of $\mathbb{C} \setminus \Gamma$ are called *basic regions* for f . The following is a corollary of Proposition 3.1.

Corollary 3.3. *There are finitely many basic regions for fixed rays.*

Proof. As tracts do not accumulate in any compact set, there are only finitely many tracts intersecting D . As preimages of δ do not accumulate in any compact set either, there are finitely many fundamental domains intersecting D , hence by Proposition 3.1 all but finitely many fixed rays must land at repelling fixed points in the interior of their respective fundamental domains. Therefore, only the finitely many remaining ones are free to land together with other rays. \square

This concludes the proof of Proposition A. The next two lemmas are interesting but will not be used later on.

Lemma 3.4 (Crossing tracts). *A basic region cannot be fully contained inside a tract.*

Proof. This proof uses notation from Subsections 2.2 and 2.4. If a basic region is fully contained inside some tract T_α , there exist at least two fixed rays g^1 and g^2 landing at a common fixed point z which are completely contained in T_α . Let $\{z^k\}$ be the fiber of z under the exponential map, with the convention that $z^k \in \tilde{S}_k$. Because z is fixed, there is a unique

$z^* \in \{z^k\}$ which is fixed. Let \tilde{g}^1 and \tilde{g}^2 be the lifts of g^1, g^2 landing at z^* . It follows that these curves must be fixed ray tails for \tilde{f} : indeed, the unique lifts of g^1, g^2 which are fixed are both forced to land at the unique fixed point. This gives two fixed ray tails for \tilde{f} which are contained in the same tract, contradicting Theorem 2.7. \square

Lemma 3.5. *Fixed ray pairs separate the set of singular values.*

Proof. If a ray pair does not separate the set of singular values, the disc D can be redefined as to not intersect the ray pair. Then, the original ray pair is contained in $\mathbb{C} \setminus \overline{D}$, hence its preimages (which includes the rays themselves) are fully contained inside tracts, contradicting Lemma 3.4. \square

4 Tools: index and homotopies

To count the number of fixed points of f inside a basic region we will use the Argument Principle. The following are preliminary concepts about index of arcs and its invariance under homotopy. For a general theory about index see for example [Why].

In this section a *curve* γ is a continuous map $\gamma : [a, b] \rightarrow \mathbb{C}$, with $a, b \in \mathbb{R}$, $a \leq b$. A curve is closed if $\gamma(a) = \gamma(b)$. An *arc* is a non-closed curve. Abusing notation, the symbol γ may denote both the function $\gamma(t)$ or the set $\gamma[a, b]$. If γ is closed and injective it is called a *Jordan curve*. For a curve γ as above and a point $P \in \mathbb{C} \setminus \gamma$ there exists a continuous branch $u(t)$ of the argument of $u(t) - P$. The *index* of γ with respect to P is then defined as

$$\text{Ind}(\gamma, P) := \frac{1}{2\pi} (u(b) - u(a)),$$

a real number independent of the chosen branch of the argument. If γ is rectifiable then the index can be expressed as the integral

$$\text{Ind}(\gamma, P) = \text{Re} \frac{1}{2\pi i} \int_{\gamma} \frac{1}{z - P} dz.$$

By definition, if γ is the union of consecutive arcs γ_i , then $\text{Ind}(\gamma, P) = \sum_i \text{Ind}(\gamma_i, P)$.

If γ is closed, $\text{Ind}(\gamma, P)$ is the winding number of γ with respect to P and it is an integer. In this case it can be used to count the number of zeros of a holomorphic map inside a Jordan curve.

Theorem 4.1 (Argument principle). *Let γ be a Jordan curve bounding a region Ω . Let f be holomorphic in a neighborhood of $\overline{\Omega}$ such that $f(z) \neq 0$ for all points $z \in \gamma$. Let $Z(f)$ be the set of zeros of f . Then*

$$\text{Ind}(f(\gamma), 0) = \#(Z(f) \cap \Omega)$$

counted with multiplicity.

Remark 4.2. The Argument principle is most often stated for rectifiable curves to be able to use the integral expression of the index. But every Jordan curve γ can be approximated by a piecewise linear Jordan curve which bounds the same zeros of f and such that its image has the same index as $f(\gamma)$ (see e.g. [Why, Sect V.3]).

Given two curves $\gamma, \sigma : [a, b] \rightarrow \mathbb{C}$, we denote by $\sigma - \gamma : [a, b] \rightarrow \mathbb{C}$ the curve defined by $\sigma(t) - \gamma(t)$. The following is a corollary of the Argument Principle, and will be our main tool.

Corollary 4.3 (Counting fixed points). *Let γ be a Jordan curve bounding a region Ω . Let f be holomorphic in a neighborhood of $\overline{\Omega}$ such that $f(z) \neq z$ for all points $z \in \gamma$. Let $\text{Fixed}(f)$ be the set of fixed points of f . Then*

$$\text{Ind}(f(\gamma) - \gamma, 0) = \#(\text{Fixed}(f) \cap \Omega)$$

counted with multiplicity.

In the remaining sections, we shall apply this corollary repeatedly, and we will do it piecewise, i.e. breaking the Jordan curve in a finite number of arcs and computing the contribution of each piece to the total index. Hence given two curves $\gamma(t)$ and $\sigma(t) := f(\gamma(t))$, not necessarily closed, we will need to compute the index of the *subtraction curve* $\Gamma(t) := \sigma(t) - \gamma(t)$ with respect to 0, in a variety of different situations. This is a more general problem in which the role of f is substituted by the relation between the respective parametrizations of σ and γ . In what follows, we collect some lemmas which give this index in terms of the relative positions and parametrizations of σ and γ . They are based on the invariance of the index under homotopies satisfying certain conditions. We first recall the definition of (relative) homotopy.

Definition 4.4 (Relative homotopies). Let $\gamma, \hat{\gamma} : [a, b] \rightarrow X \subset \mathbb{C}$ be two curves. We say that γ and $\hat{\gamma}$ are *homotopic in X* if there exists a continuous map

$$\begin{aligned} H : [a, b] \times [0, 1] &\longrightarrow X \\ (t, s) &\longmapsto H(t, s) =: \gamma_s(t) \end{aligned}$$

such that $\gamma_0(t) = \gamma(t)$ and $\gamma_1(t) = \hat{\gamma}(t)$ for all $t \in [a, b]$. If $\gamma_s(t_0) = P_0$ for some $t_0 \in [a, b]$, some $P_0 \in X$ and for all $s \in [0, 1]$, we say that γ and $\hat{\gamma}$ are homotopic in X relative to P_0 , and write $\gamma \sim \hat{\gamma} \text{ rel } \{P_0\}$ in X . If the homotopy is relative to a set $A \subset X$, it is understood that it is relative to each $P_0 \in A$. In particular, a curve is *homotopic in X to a point P_0* if it can be continuously deformed *in X* to the constant curve equal to P_0 .

Lemma 4.5 (First Homotopy Lemma).

- (a) *Fix $P \in \mathbb{C}$ and let $\gamma, \hat{\gamma} : [0, 1] \rightarrow \mathbb{C} \setminus \{P\}$ be two curves with common endpoints P_0, P_1 . Suppose that γ and $\hat{\gamma}$ are homotopic in $\mathbb{C} \setminus \{P\}$, relative to P_0 and P_1 . Then*

$$\text{Ind}(\gamma, P) = \text{Ind}(\hat{\gamma}, P).$$

- (b) *Let $\gamma, \hat{\gamma}, \sigma, \hat{\sigma} : [0, 1] \rightarrow \mathbb{C}$ be curves such that $\gamma \sim \hat{\gamma} \text{ rel } \{\gamma(0), \gamma(1)\}$ and $\sigma \sim \hat{\sigma} \text{ rel } \{\sigma(0), \sigma(1)\}$, both homotopies in \mathbb{C} . Assume further that $\sigma_s(t) - \gamma_s(t) \neq 0$ for all $s, t \in [0, 1]$. Then*

$$\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\hat{\sigma} - \hat{\gamma}, 0).$$

See Figure 6. The proof is left as an exercise. Part (b) follows from (a) using that the subtraction of the two homotopies gives a homotopy between $\Gamma(t) := \sigma(t) - \gamma(t)$ and $\widehat{\Gamma}(t) := \widehat{\gamma}(t) - \widehat{\sigma}(t)$ in $\mathbb{C} \setminus \{0\}$ relative to their common endpoints.

We shall use part (b) repeatedly throughout this section and inside the proof of the Main Theorem. Informally, it says that we can deform each curve separately as long as we do not create zeros of the subtraction curves Γ_s at any step. Observe that a continuous series of reparametrizations of a curve is a homotopy. For single curves, this has no effect in the index. But in the setup above, reparametrizing one of the curves does affect Γ , since it changes how the points in γ and σ correspond.




Figure 6: Setup of the First Homotopy Lemma part (a) (left) and part (b) (right). In this particular example, because of the way that γ and σ correspond, we may deform σ and γ (continuous) into $\widehat{\sigma}$ and $\widehat{\gamma}$ (dashed) without creating any zero of $\sigma_s(t) - \gamma_s(t)$ in the way. This means that $\text{Ind}(\sigma_s - \gamma_s, 0)$ is constant for all s . The dotted curves denote $\gamma_s(0.5)$ and $\sigma_s(0.5)$ respectively.

With the first Lemma as a tool we prove the following. See Figures 7 and 8.

Lemma 4.6 (Second Homotopy Lemma). *Let $\gamma, \sigma : [0, 1] \rightarrow \mathbb{C}$ be two curves such that $\sigma(0) = \sigma(1) = P \in \mathbb{C}$. Suppose one of the following occurs:*

- (a) $\gamma \cap \sigma = \emptyset$; or
- (b) $\gamma \cap \sigma \neq \emptyset$ but
 - (i) $\gamma(t) \neq \sigma(t)$ for all $t \in [0, 1]$, and $P \notin \gamma$;
 - (ii) there exists a connected component U of $\mathbb{C} \setminus \sigma$, such that $\gamma(0), \gamma(1) \in U$ and $\gamma(0, 1) \subset \overline{U}$;
 - (iii) if $z \in \partial U$ has more than one access from U , then $z \neq \gamma(t)$ for any $t \in [0, 1]$.

Then

$$\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\gamma, P) + N,$$

where $N = \text{Ind}(\sigma, z)$ for all $z \in \gamma \setminus \sigma$.




Figure 7: Setup of the Second Homotopy Lemma part (a) with $N = 0$ (left) and $N = 2$ (right).




Figure 8: Setup of the Second Homotopy Lemma part (b) with $N = 0$ (left) and $N = 2$ (right). The component U is shadowed. The broken lines show the curve γ_1 (to be constructed in the proof) homotopic to γ and not intersecting σ .

Proof. Case (a): Since γ and σ do not intersect, all points in γ must be in the same component U of $\mathbb{C} \setminus \sigma$. Hence the index N is well defined for all $z \in \gamma$ and does not depend on z .

Observe that the curve $\sigma(t) - \gamma(t)$ starts with the vector $P - \gamma(0)$ and ends with the vector $P - \gamma(1)$. Up to an integer, the difference between the arguments of these two vectors is actually the index of γ with respect to P , but also the index of $\sigma - \gamma$ with respect to 0 . Hence $\text{Ind}(\sigma - \gamma, 0) - \text{Ind}(\gamma, P) \in \mathbb{Z}$. We will prove that this integer equals N by proving that for every $\delta > 0$ we can find $\varepsilon > 0$ and a curve Γ_ε such that $\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\Gamma_\varepsilon, 0)$ and

$$|\text{Ind}(\Gamma_\varepsilon, 0) - \text{Ind}(\gamma, P) - N| < \delta.$$

The result then follows since the index must be an integer.

For every $\varepsilon > 0$, we first make a continuous series of reparametrizations γ_s^ε of γ , $s \in [0, 1]$, so that the final curve γ_1^ε is constant and equal to $\gamma(0)$, for all $t \in [0, 1 - \varepsilon]$ and then “runs” to $\gamma(1)$ in the remaining ε time. An explicit homotopy is for example

$$\gamma_s^\varepsilon(t) := \begin{cases} \gamma(0) & \text{if } t \in [0, s(1 - \varepsilon)] \\ \gamma\left(\frac{t - s(1 - \varepsilon)}{1 - s(1 - \varepsilon)}\right) & \text{if } t \in [s(1 - \varepsilon), 1]. \end{cases}$$

Now define $\Gamma_\varepsilon := \sigma - \gamma_1^\varepsilon$. Since the curves are disjoint at every step, Γ_ε is homotopic to $\sigma - \gamma$ (relative to its endpoints) in $\mathbb{C} \setminus \{0\}$. Observe that $\Gamma_\varepsilon(t) = \sigma(t) - \gamma(0)$ for all $t \in [0, 1 - \varepsilon]$, hence the contribution to the index during this parameter interval is arbitrarily close to N if ε is small enough. On the other hand, for $t \in [1 - \varepsilon, 1]$ we have $\Gamma_\varepsilon(t) \simeq P - \gamma(t)$, so the index contribution is arbitrarily close to $\text{Ind}(\gamma, P)$ for ε sufficiently small. It follows that $\text{Ind}(\Gamma_\varepsilon)$ is arbitrarily close to $\text{Ind}(\gamma, P) + N$ as we wanted to show.

Case (b): Observe that N is also well defined in this case for the points $z \in \gamma \setminus \sigma$. In this setting the trick of part (a) does not necessarily work since it is not clear that by reparametrizing γ we do not create zeros of the subtraction curve. The idea is to make a homotopy which moves γ away from σ . More precisely we will define a homotopy from γ to a curve γ_1 in such a way that $\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\sigma - \gamma_1, 0)$ and $\gamma_1 \cap \sigma = \emptyset$ so that the result follows from case (a).

To that end, observe that U is a topological disc (in $\widehat{\mathbb{C}}$) with locally connected boundary, since σ is connected and locally connected (the image of a closed interval is locally connected by Hahn-Mazurkiewicz Theorem, and each component of $\mathbb{C} \setminus \sigma$ also is by Torhorst's Theorem [Why]). Let $\varphi : \mathbb{D} \rightarrow \overline{U}$ be a Riemann map extended to the boundary and set $\widehat{\gamma}(t) := \varphi^{-1}(\gamma(t)) \in \mathbb{D}$. Observe that $\varphi^{-1}(\sigma \cap \gamma)$ is contained in $\{|z| = 1\}$. Fix $0 < \rho < 1$ such that $\widehat{\gamma}(0), \widehat{\gamma}(1) \in \mathbb{D}_\rho$ and let $\varepsilon < 1 - \rho$. For $r, s \in [0, 1]$ let l_s be the piecewise linear map which is constant on $[0, \rho]$ and affine from $[\rho, 1]$ onto $[\rho, 1 - s\varepsilon]$. Observe that l_s depends continuously on s . Now define $h_s : \mathbb{D} \rightarrow \mathbb{D}_{1-s\varepsilon}$ as $h_s(re^{i\theta}) = l_s(r)e^{i\theta}$. Then h_s is an angle-preserving homeomorphism which is the identity on \mathbb{D}_ρ , and sends the annulus $\mathbb{D} \setminus \mathbb{D}_\rho$ to the annulus $\mathbb{D}_{1-s\varepsilon} \setminus \mathbb{D}_\rho$, satisfying that $h_0 = \text{Id}$ on \mathbb{D} . Hence the map

$$\widehat{H}(t, s) := \widehat{\gamma}_s(t) := h_s(\widehat{\gamma}(t))$$

is continuous in both variables and a homotopy between $\widehat{\gamma}$ and $\widehat{\gamma}_1$ relative to their endpoints $\varphi(\gamma(0))$ and $\varphi(\gamma(1))$. It follows from (iii) that

$$H(t, s) := \gamma_s(t) := \varphi(\widehat{\gamma}_s(t))$$

defines a homotopy between γ and γ_1 relative to its endpoints. It is clear that $\sigma(t) - \gamma_s(t) \neq 0$ for all $s, t \in [0, 1]$ hence, by Lemma 4.5, we deduce that $\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\sigma - \gamma_1, 0)$. Since σ is disjoint from γ_1 , the result then follows from case (a). \square

5 Global counting: proof of Theorem B

Recall that, given a fundamental domain F and its fixed dynamic ray g_F as given by the Structural Lemma 2.3, we say that g_F *lands alone* if it lands, and no other fixed ray has the same landing point. A ray may land alone at a repelling fixed point or at a parabolic fixed point. By Proposition 3.1, there are only finitely many fundamental domains whose fixed rays do not land alone at repelling fixed points.

Definition 5.1 (Full and complete collections). A collection \mathcal{F}_α of fundamental domains in T_α is *full* if for any $F, F' \in \mathcal{F}_\alpha$, the collection contains any other $F \in T_\alpha$ in between F and F' with respect to the order of the fundamental domains in T_α . A collection \mathcal{F} of fundamental

domains in \mathcal{T} is *full* if its restriction to each one of the tracts is full. We say that \mathcal{F} is *complete* if it contains all fundamental domains whose fixed ray does not land alone at a repelling fixed point, and it also includes all fundamental domains which intersect D .

Informally, a collection is full if it has no gaps (in each tract). Theorem B is an easy consequence of the following theorem.

Theorem 5.2 (Global counting). *Let $f \in \widehat{\mathcal{B}}$ and assume that all fixed rays land. Let \mathcal{F} be a finite collection of fundamental domains which is full and complete. Let $N = \#\mathcal{F}$ and \mathcal{G} be the collection of N fixed rays which are asymptotically contained in some $F \in \mathcal{F}$. Then there are exactly $N + 1$ fixed points counted with multiplicity, which are either landing points of some $g \in \mathcal{G}$, or interior fixed points.*

Let us first show how Theorem B follows from Theorem 5.2.

Proof of Theorem B. Let G be the finite collection of L fixed rays given in Theorem B, which includes all rays which do not land alone at repelling fixed points. Let \mathcal{F}' be the finite collection of L fundamental domains in which the rays in G are asymptotically contained. By adding a finite number of fundamental domains to \mathcal{F}' , (whose fixed rays must land alone at repelling fixed points), we obtain a new finite collection $\mathcal{F} \supset \mathcal{F}'$ which is complete and full (informally, we first add all fundamental domains which intersect D and are not already in \mathcal{F}' , and then "fill in" the gaps in between non-consecutive domains). Set $N = \#\mathcal{F}$, and let $\mathcal{G} \supset G$ be the collection of $N \geq L$ rays associated to \mathcal{F} . By Theorem 5.2 there exist $N + 1$ fixed points counted by multiplicity which are either landing points of some $g \in \mathcal{G}$, or interior fixed points. Now note that the $N - L$ rays in $\mathcal{G} \setminus G$ are necessarily fixed rays which land alone at repelling (hence simple) fixed points. Therefore there are exactly $N - L$ simple fixed points which are landing points of these $N - L$ rays. Thus the remaining $L + 1$ fixed points (counted with multiplicity) are either landing points of a ray $g \in G$ or interior fixed points. \square

Before proving Theorem 5.2 we would like to emphasize the following simple but rather surprising corollary.

Corollary 5.3. *Let $f \in \widehat{\mathcal{B}}$ and assume that all fixed rays land. If there is only one basic region (with respect to the set of fixed rays), then there is exactly one interior fixed point or virtual fixed point.*

Proof. If there is only one basic region then there are no fixed ray pairs, that is, all fixed rays land alone either at repelling fixed points or parabolic ones. There are only a finite number of the latter by Proposition A. Let G be *any* collection of N rays which includes them. By Theorem B there are $N + 1$ fixed points counted with multiplicity which are either landing points of some $g \in G$ or interior fixed points. But all rays land alone; so, either all the landing fixed points are repelling and there is exactly one fixed point left which must be an interior fixed point, or one (and only one) of the landing points is parabolic of multiplicity two and there is one (and only one) virtual fixed point. \square

This means for example, that even if f may have infinitely many singular values, if all fixed rays land alone, there is at most one non-repelling fixed point.

The remaining of the section is dedicated to prove Theorem 5.2.

Proof of Theorem 5.2. Let \mathcal{F} , \mathcal{G} and N be as in the statement. Since the collections are finite, their members intersect a finite number of tracts say, $\{T_\alpha\}_{1 \leq \alpha \leq M}$, where the tracts are labeled consecutively respecting the cyclic order. From now on the index α will be taken modulo M . Let $\mathcal{F}_\alpha := \mathcal{F} \cap T_\alpha$, and $N_\alpha := \#\mathcal{F}_\alpha$.

Claim 5.4. *Without loss of generality we may assume that*

- (a) δ does not intersect the set of fixed rays and
- (b) T_α intersects D for every $1 \leq \alpha \leq M$.

Proof. In fact we will show that (a) and (b) are satisfied up to considering a full and complete finite family \mathcal{F}' of fundamental domains, which is obtained by adding to \mathcal{F} finitely many fundamental domains (whose fixed ray must land alone at a repelling fixed point). Proving Theorem 5.2 for \mathcal{F}' then implies Theorem 5.2 for \mathcal{F} .

Observe that rays are intrinsic to the function f , i.e., they do not depend on the definitions of D or δ . Tracts and fundamental domains do. By Lemma 2.12 there exists a sufficiently large disc $D' \supset D$ such that the set of fixed rays is contained in $D' \cup \mathcal{T}$, where \mathcal{T} are the tracts defined by D . Also, for any tract T_α which contains at least one fundamental domain $F_\alpha \in \mathcal{F}$, let z_α be the landing point of the fixed ray associated to F_α ; up to enlarging D' , we can assume that it contains all such z_α . Let δ' be the subset of δ in $\mathbb{C} \setminus D'$ connecting D' with infinity. By construction δ' does not intersect any fixed ray. The tracts defined as preimages of $\mathbb{C} \setminus D'$ are smaller than the previous ones, however, the set of fixed rays does not change. Also, the new fundamental domains are asymptotically the same as the old fundamental domains, since $\delta' = \delta$ outside a compact set. By Lemma 3.1, and for all $\alpha = 1 \dots M$, the new fundamental domains F'_α which are contained in F_α all intersect D' , since their fixed ray lands in D' .

The enlarged disk D' intersects possibly more fundamental domains than the ones which were originally in \mathcal{F} , so \mathcal{F} is no longer a full and complete collection with respect to the new disc. However it can be turned into a full and complete collection \mathcal{F}' (which satisfies property (b) by construction), by adding finitely many fundamental domains, whose fixed rays all land alone at repelling fixed points since \mathcal{F} was originally complete. \square

Let C_R denote the circle or radius R . Choose R large enough so that part (b) of the Structural Lemma 2.3 is satisfied for the family of fundamental domains \mathcal{F} , and such that all fixed rays associated to fundamental domains in \mathcal{F} land inside the disc D_R .

Let P be the first intersection of δ with C_R when moving from D towards infinity. Call δ_P the arc in δ connecting P to D . Let δ_α^\pm be the two preimages of δ_P which are contained in T_α , so that each δ_α^\pm is on the boundary between a fundamental domain in \mathcal{F}_α and a fundamental domain which is not in \mathcal{F}_α (these curves are well defined because \mathcal{F} is full). Label the δ_α^\pm so as to respect the cyclic order in the sense that δ_α^- comes before δ_α^+ and after $\delta_{\alpha-1}^+$. Each

δ_α^\pm intersects the preimage of C_R in a point P_α^\pm which is a preimage of P , and ends at the boundary of T_α . Let r_α be the arc in the preimage of C_R connecting P_α^- and P_α^+ inside T_α and such that $f(r_\alpha)$ covers C_R N_α times. As usual, give C_R and any other Jordan curve in this section a counterclockwise orientation; this induces an orientation on the arcs r_α (see Figure 9 for an illustration of a possible setup).

Finally, let γ_α be Jordan arcs in $\mathbb{C} \setminus (\overline{\mathcal{T} \cup D})$ connecting $\delta_{\alpha-1}^+$ with δ_α^- , which do not intersect the set of fixed rays (this can be done, because only finitely many fixed rays intersect $\mathbb{C} \setminus \overline{\mathcal{T}}$ by Lemma 3.1). Observe that $\bigcup T_\alpha \cup D$ divides the plane into finitely many components, only one of which contains δ . In view of Claim 5.4, all tracts T_α intersect D , so one and exactly one of the curves γ_α intersects δ . Choose that specific γ_α so as to intersect δ only once, and to do so in the set $\{|z| > R\}$. Note that the image of each γ_α is fully contained in D .

Call Γ_α the arcs $\delta_{\alpha-1}^+ \cup \gamma_\alpha \cup \delta_\alpha^-$ connecting $P_{\alpha-1}^+$ to P_α^- . We will count the number of fixed points of f inside the region \widehat{U} enclosed by the Jordan curve

$$\Gamma = \bigcup_{\alpha \leq M} (\Gamma_\alpha \cup r_\alpha).$$

Observe that, for R large enough, all fixed points which are landing points of the fixed rays associated to the fundamental domains in \mathcal{F}_α have to be contained in \widehat{U} . Also, because there are no fixed points in $\mathbb{C} \setminus (\overline{D \cup \mathcal{T}})$, the final count does not depend on the choice of γ_α . We will also see that the result does not depend on C_R (and hence on the r_α) as long as C_R satisfies the Structural Lemma 2.3 (see Figure 9).

We calculate $\text{Ind}(f(\Gamma) - \Gamma, 0)$, by calculating it piece by piece on the r_α and on the Γ_α . Observe that by construction there are no fixed points on Γ . Let us first calculate $\text{Ind}(f(r_\alpha) - r_\alpha, 0)$. Observe that $f(r_\alpha)$ is the circle C_R covered N_α times, starting and ending at P . Then by the Second homotopy Lemma 4.6, part (a),

$$\text{Ind}(f(r_\alpha) - r_\alpha, 0) = N_\alpha + \text{Ind}(r_\alpha, P).$$

Now let us calculate $\text{Ind}(f(\Gamma_\alpha) - \Gamma_\alpha, 0)$. The image of each Γ_α is a curve starting at P , moving along δ_P , staying inside \overline{D} and then moving back to P along δ_P .

Observe that $\Gamma_\alpha \cap f(\Gamma_\alpha) = \emptyset$. Then, by the Second Homotopy Lemma 4.6, part (b), with $N = 0$ we have

$$\text{Ind}(f(\Gamma_\alpha) - \Gamma_\alpha, 0) = \text{Ind}(\Gamma_\alpha, P).$$

Summing up, we obtain

$$\text{Ind}(f(\Gamma) - \Gamma, 0) = \sum_{\alpha} \left(\text{Ind}(r_\alpha, P) + N_\alpha + \text{Ind}(\Gamma_\alpha, P) \right) = N + \text{Ind}(\Gamma, P) = N + 1.$$

The last equality follows from the fact that Γ is a Jordan curve (oriented counterclockwise) and P is contained in the bounded connected component of $\mathbb{C} \setminus \Gamma$. Hence, there are exactly $N + 1$ fixed points in \widehat{U} counted with multiplicity. \square




Figure 9: Sketch of a setup for the global counting. The curve Γ is the boundary of the shaded region \widehat{U} . Its image consists of the big circle C_R , together with δ_P and the images of the curves γ_i which are inside D . In this example the Argument Principle gives 8 fixed points inside the shaded region, counted with multiplicity. Since there are only 7 fixed rays landing in \widehat{U} , at least one of the fixed points must be an interior or virtual fixed point.

6 Local Counting. Proof of the Main Theorem

If there exists only one basic region, the Main Theorem reduces to Corollary 5.3. So from now on we assume that there are at least two basic regions.

Let us consider a basic region V and show that it contains exactly one interior fixed point or a virtual fixed point. For the purpose of the proof in this section, we redefine V to contain also the fixed rays landing alone (at repelling or parabolic fixed points) that were part of ∂V . In other words, ∂V now consists exclusively of ray pairs. Without loss of generality, since there are at least two basic regions, we can also assume that $\delta \cap V = \emptyset$ (see Lemma 2.1, Lemma 3.4 and Claim 5.4). In fact, redefining δ and/or D will redefine fundamental domains, but not fixed rays nor V itself.

Because of the presence of fixed rays and fixed points in ∂V , the counting is more involved than it was in the last section. Notice however that the only fixed points in ∂V are landing points of fixed rays, which by the Snail lemma [Mi, Lemma 16.2] must be repelling or parabolic.

The strategy of the proof is as follows. We first consider the case where V contains no virtual fixed point. This implies that every boundary fixed point is repelling or, if it

is parabolic, both boundary rays land at it from the same repelling petal (see the precise definition below). We then define a new region \widehat{V} , a bounded modification of V , which includes all the repelling fixed points which were previously on ∂V and excludes all the parabolic ones. After defining $\Gamma_{\widehat{V}} := \partial\widehat{V}$, now a fixed point free Jordan curve, we will apply the argument principle to $f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}$ to count the number of fixed points in \widehat{V} . We shall show that \widehat{V} contains as many fixed points as the number of rays in \widehat{V} landing alone, plus the number of repelling fixed points which were previously in ∂V , plus one extra fixed point which must therefore be interior. Note that this extra fixed point cannot be attributed to a fixed point of multiplicity two since we assumed that V had no virtual fixed points to start with and we left out all parabolic points which were previously on ∂V .

To that end, we shall compute $\text{Ind}(f(\widehat{\Gamma}) - \widehat{\Gamma}, 0)$ piecewise, as in the section above, using the Homotopy Lemmas in Section 4, but also doing some homotopies *in situ*. It is worth noticing that homotopies are used exclusively to compute the indices of certain arcs. They do not further modify the region \widehat{V} or the map f .

Finally we deal with the remaining regions which contain virtual fixed points. This will be done by a global counting argument.

Modification of V near the fixed points on the boundary

By definition basic regions are open. Suppose $z_0 \in \partial V$ is a repelling or parabolic fixed point. We slightly modify the boundary of V near z_0 making use of the local dynamics around z_0 . Here we describe the modification near every fixed point on ∂V ; in the proof of the Main Theorem we will apply all the modifications together. In these lemmas V' denotes the modification of V .

Lemma 6.1 (Repelling fixed points in ∂V). *Let z_0 be a repelling fixed point on ∂V . Then, a small arc on ∂V containing z_0 can be replaced by an arc ζ with the same endpoints such that:*

- *the resulting region $V' \supset V$ satisfies $\text{Fixed}(f) \cap V' = (\text{Fixed}(f) \cap V) \cup \{z_0\}$;*
- *f has no fixed points on ζ ;*
- *$f(\zeta)$ is contained in $\mathbb{C} \setminus V'$.*

Proof. Let D_ε be a small disc centered at z_0 , with ε small enough so that $f(D_\varepsilon)$ is a topological disc containing $\overline{D_\varepsilon}$. See Figure 10. We define V' as the “filled” region $V \cup D_\varepsilon$, that is, the region $V \cup D_\varepsilon$ union the (finite number of) bounded connected components of its complement (which are due to the possible wiggling of the rays in and out D_ε). By choosing ε small enough, we can make sure that we added no extra fixed points to V' other than z_0 . We then define ζ as the arc in $\partial V'$ starting and ending at ∂D_ε such that $\partial V' \setminus \zeta \subset \partial V$. More precisely, ζ joins the two points x_1 and x_2 in ∂D_ε which correspond to the first intersection (when coming from infinity) of the fixed rays in ∂V landing at z_0 , with ∂D_ε . Observe that if these rays do not wiggle outside of D_ε after entering it for the first time, then $\zeta \subset \partial D_\varepsilon$. Otherwise, ζ also contains parts of the rays.

By construction $f(\zeta)$ has no fixed points and is contained in $\mathbb{C} \setminus V'$. Indeed the part of ζ on ∂D_ε is mapped to $\mathbb{C} \setminus \overline{V'}$ while the part of ζ which is on the rays is mapped to the rays (at a higher potential than the one of x_1 or x_2), hence to $\partial V'$. \square

We now consider the case of parabolic fixed points. For the following description in more detail see for example [Mi, §10]. Let z_0 be a parabolic fixed point of multiplicity $m + 1$. On any simply connected neighborhood of z_0 disjoint from the set of singular values, there is a well defined branch φ of f^{-1} which fixes z_0 . In this setup, there exist m *attracting vectors* such that if an orbit converges to z_0 it does so in a direction asymptotically tangent to one of them. Likewise, there exist m *repelling vectors*, defined as attracting vectors for φ . Attracting and repelling vectors alternate. An *attracting petal* at z_0 is defined as any open simply connected set W containing z_0 on its boundary, such that $f(\overline{W}) \subset W \cup \{z_0\}$. Likewise, a *repelling petal* is an attracting petal for φ . Every attracting (resp. repelling) petal is associated to an attracting (resp. repelling) vector. Every virtual fixed point (or immediate attracting basin) attached to z_0 is associated to a unique attracting vector and contains all attracting petals also associated to the same vector.

Lemma 6.2 (Parabolic fixed points in ∂V). *Let z_0 be a parabolic fixed point on ∂V , such that no virtual fixed point associated to z_0 is contained in V . Then, a small arc on ∂V containing z_0 can be replaced by an arc ζ with the same endpoints such that:*

- *the resulting new region $V' \subset V$ satisfies $\text{Fixed}(f) \cap V' = \text{Fixed}(f) \cap V$;*
- *f has no fixed points on ζ ;*
- *$f(\zeta)$ is contained in $\overline{V'}$.*




Figure 10: Left: The enlargement of V near a repelling fixed point on its boundary. The region V' includes the lighter shaded region V as well as the darker shaded parts. Right: The shrinking of V near a parabolic fixed point in its boundary.

Proof. Since there is no virtual fixed point associated to z_0 in V , there exists an arbitrarily small repelling petal W attached to z_0 which eventually contains both rays landing at z_0 on ∂V . See Figure 10. Consider the region V' to be the unbounded connected component of $V \setminus \overline{W}$. We define $\zeta \subset \partial V'$ as the curve starting and ending in ∂W , not going through z_0 , such that $\partial V' \setminus \zeta \subset \partial V$. Equivalently ζ joins the first intersection points between the rays and ∂W . If the rays do not wiggle outside of W once they enter it, then $\zeta \subset \partial W$, but otherwise ζ contains part of the rays. By construction, the new region V' does not contain z_0 and f has no fixed points on ζ . Finally, $f(\zeta)$ is contained in V' for the part of ζ in ∂W , and $f(\zeta)$ is contained in $\partial V'$ for the part of ζ in the rays. Again, by shrinking W if necessary, we can make sure we did not remove any fixed point from V by cutting away the bounded components of $V' \setminus W$. \square

Proof of the Main Theorem

The setup for the proof of the Main Theorem is very similar to the setup in Section 5, so we will use the notation introduced there. Let \mathcal{F} be a finite full and complete collection of fundamental domains (see Definition 5.1). Let $P, \Gamma_\alpha, r_\alpha$ be as in the proof of Theorem 5.2, and \widehat{U} be the region bounded by $\bigcup(r_\alpha \cup \Gamma_\alpha)$. Let V be a basic region for f . Recall that we only have to consider the case in which there are at least two basic regions, so that without loss of generality $V \cap \delta = \emptyset$. Assume first that there are no virtual fixed points in V . Let $\widehat{V} = V' \cap \widehat{U}$, where V' is V after being modified near the boundary fixed points as in Lemmas 6.1 and 6.2. Recall that when modifying the region, repelling fixed points on ∂V end up inside V' , while parabolic fixed points do not. Without loss of generality, R is large enough so that all fixed rays associated to the fundamental domains in $\mathcal{F} \cap V$ have their landing points in \widehat{V} . Let $\Gamma_{\widehat{V}}$ be the boundary of \widehat{V} .

The difference with respect to the global counting in Section 5 is that we additionally have to calculate the contributions to $\text{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}, 0)$ made by the (pieces of) fixed ray pairs in $\Gamma_{\widehat{V}}$. Observe that there is one and exactly one fixed ray pair on $\Gamma_{\widehat{V}}$ separating δ from \widehat{V} . Let $\{P_i\}_{i=1}^k$ be the collection of k preimages of P which are contained in $\Gamma_{\widehat{V}}$, labeled so as to respect the order in which they are encountered when moving along $\Gamma_{\widehat{V}}$ counterclockwise. Also, let s_i be the arc along $\Gamma_{\widehat{V}}$ joining two consecutive points P_i and P_{i+1} , where indices are taken mod k (see Figure 11).

The arcs s_i can be of four types:

- (0) s_i intersects the ray pair separating δ from \widehat{V} ;
- (1) s_i intersects any other ray pair;
- (2) $s_i \subset r_\alpha$ for some r_α ;
- (3) $s_i = \Gamma_\alpha$ for some α .

We only have to compute $\text{Ind}(f(s_i) - s_i, 0)$ when s_i is of type (0) and (1), as the other two cases are already covered in the previous section. We calculate this in two separate claims. For each s_i of type (0) and (1), let $n_i = 1$ if the ray pair originally landed at a repelling fixed point, and $n_i = 0$ if the ray pair originally landed at a parabolic fixed point.




Figure 11: A simple example of counting for the modified basic region \widehat{V} shown (shaded) in the picture. The original region V had two repelling fixed points on its boundary which are now part of \widehat{V} . Arcs of type (0) and (1) are colored in blue; arcs of type (2) are colored in green; arcs of type (3) are in red. Each arc has the same color as its image. In this example $\text{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}) = 7$.

Claim 6.3. *If s_i is of type (0), the counting gives $\text{Ind}(f(s_i) - s_i, 0) = 1 + n_i + \text{Ind}(s_i, P)$.*

Proof. Let s_i be the arc of type (0), g_1 and g_2 be respectively the first and the second ray in s_i that we encounter when moving along $\Gamma_{\widehat{V}}$ counterclockwise. Let ζ be as in Lemma 6.1 or 6.2 depending on whether the original fixed point was repelling ($n_i = 1$) or parabolic ($n_i = 0$). Then $f(s_i)$ starts at P , goes along C_R counterclockwise until meeting g_1 , continues on g_1 until hitting ζ , then along $f(\zeta)$, then moves back along the ray g_2 and continues along C_R until it meets P again. See Figure 12. Note that $f(s_i)$ divides the plane into exactly three connected components. Denote by U_1 the bounded component with P on the boundary and by U_2 the bounded component which contains the points P_i and P_{i+1} . Observe that $\text{Ind}(f(s_i), z) = 1$ if $z \in U_1$ and equals 2 if $z \in U_2$.

Let us first suppose that $n_i = 1$. Then by Lemma 6.1 $f(\zeta)$, and hence $f(s_i)$, is contained in $\mathbb{C} \setminus \widehat{V}$, which is closed. Hence s_i , and in fact the whole \widehat{V} , is contained in $\overline{U_2}$, so that $\text{Ind}(f(s_i), z) = 2$ for all $z \in s_i \setminus f(s_i)$. By the Second Homotopy Lemma 4.6 part (b), with $\gamma = s_i$ and $\sigma = f(s_i)$ we obtain that $\text{Ind}(f(s_i) - s_i, 0) = 2 + \text{Ind}(s_i, P)$.

Suppose now that $n_i = 0$. Then, by Lemma 6.2, $f(\zeta)$ is contained in the closure of \widehat{V} . This is a problem when trying to apply Lemma 4.6 since there is a part of s_i contained in $\overline{U_1}$, namely the rays and the arc ζ , while the remaining part belongs to U_2 , precisely two arcs say ξ_1 going from the endpoint P_i to the ray g_1 , and ξ_2 going from the endpoint P_{i+1} to the ray g_2 . See Figure 12. Let $\gamma = s_i$ parametrized by $[0, 1]$ and $f(s_i)$ be $\sigma(t) := f(\gamma(t))$.

We claim that there exists $\widehat{\sigma}$, a local modification of σ such that $\sigma \sim \widehat{\sigma}$ relative to P (and in fact to most of σ itself), with a homotopy satisfying the conditions of the First Homotopy

Lemma (so that $\text{Ind}(\sigma - \gamma, 0) = \text{Ind}(\widehat{\sigma} - \gamma, 0)$), and such that $\widehat{\sigma}$ and γ do satisfy the hypothesis of the Second Homotopy Lemma with $N = 1$. The curve $\widehat{\sigma}$ is constructed as follows. Let t_1 be the smallest parameter such that $\gamma(t_1) \in g_1$ and let $t'_1 > t_1$ be such that $\sigma(t'_1) = \gamma(t_1)$. Fix $\varepsilon > 0$ arbitrarily small and consider the arc $\sigma^1 = \sigma(t'_1 - \varepsilon, t'_1 + \varepsilon)$. Now consider a homotopy σ_s^1 for $s \in [0, 1]$ relative to the endpoints of σ^1 which progressively deforms the curve sliding it along $\xi_1 = \gamma[0, t_1]$ so that the end curve σ_1^1 does not intersect ξ_1 at any point. Observe that $\sigma[0, t_1]$ is very far away from σ_s^1 for all s , so that $\sigma_s^1(t) - \gamma(t) \neq 0$ for any s, t . Analogously, define a homotopy σ_s^2 in a neighborhood of ξ_2 which slides σ along ξ_2 . Finally define $\widehat{\sigma}(t) := \sigma_1^j(t)$ if $j = 1, 2$ and $t \in [t'_j - \varepsilon, t'_j + \varepsilon]$ and $\widehat{\sigma}(t) = \sigma(t)$ otherwise. If we denote by \widehat{U}_1 and \widehat{U}_2 the bounded connected components of $\mathbb{C} \setminus \widehat{\sigma}$ (which are local modifications of U_1 and U_2 respectively), we see that γ belongs to the closure of \widehat{U}_1 , so that the hypotheses of the Second Homotopy Lemma are satisfied with $N = 1$. \square




Figure 12: The computation of the index for an arc of type (0). Left: A repelling fixed point was previously in ∂V and now belongs to \widehat{V} . The arc s_i is shown in blue and belongs to \overline{U}_2 . Its image is shown in light blue. Right: A parabolic fixed point was previously in ∂V and now is not in \widehat{V} . The arc $\gamma = s_i$ is shown in blue while its image $\sigma = f(s_i)$ is shown in light blue. The dashed curves show the local modification of σ so that γ belongs to the closure of \widehat{U}_1 .

Claim 6.4. *If s_i is of type (1), the counting gives $\text{Ind}(f(s_i) - s_i, 0) = n_i + \text{Ind}(s_i, P)$.*

Proof. Let g_1 and g_2 be respectively the first and the second ray in s_i that we encounter when moving along $\Gamma_{\widehat{V}}$ counterclockwise. Let ζ be as in Lemma 6.1 or 6.2 depending on whether the original fixed point was repelling ($n_i = 1$) or parabolic ($n_i = 0$). Then $f(s_i)$ starts at P , goes along C_R until meeting g_1 , continues on g_1 until hitting ζ , then along $f(\zeta)$, then moves back along the ray g_2 and continues along C_R until meeting P again. See Figure 13. Note that in this case $f(s_i)$ is a Jordan curve which therefore divides the plane into exactly two

connected components. Denote by U_1 the bounded component of $\mathbb{C} \setminus f(s_i)$ and by U_0 the unbounded one. Then $\text{Ind}(f(s_i), z) = j$ if $z \in U_j$ for $j = 0, 1$.

As in the claim above we first suppose that $n_i = 1$ (see Figure 13). Then $s_i \subset U_1$ and by the Second Homotopy Lemma with $N = 1$ we have $\text{Ind}(f(s_i) - s_i, 0) = 1 + \text{Ind}(s_i, P)$.

Otherwise, if $n_i = 0$, we encounter the same difficulty as above: namely, part of s_i is in U_0 while the remaining part is in U_1 . We proceed analogously, by constructing a homotopy between $\sigma := f(s_i)$ and $\hat{\sigma}$ in such a way that s_i is in the closure of \widehat{U}_0 and hence $\text{Ind}(f(s_i) - s_i, 0) = \text{Ind}(s_i, P)$ by the Second Homotopy Lemma with $N = 0$. \square




Figure 13: The computation of the index for an arc of type (1). Left: A repelling fixed point was previously in ∂V and now belongs to \widehat{V} . The arc s_i is shown in blue and belongs to \overline{U}_1 . Its image is shown in light blue. Right: A parabolic fixed point was previously in ∂V and now is not in \widehat{V} . The arc $\gamma = s_i$ is shown in blue while its image $\sigma = f(s_i)$ is shown in light blue. The dashed curves show the local modification of σ so that γ belongs to the closure of \widehat{U}_0 .

So in the four cases under consideration we obtain:

$$\begin{aligned}
 \text{Ind}(f(s_i) - s_i, 0) &= 1 + n_i + \text{Ind}(s_i, P) && \text{in case (0);} \\
 \text{Ind}(f(s_i) - s_i, 0) &= n_i + \text{Ind}(s_i, P) && \text{in case (1);} \\
 \text{Ind}(f(s_i) - s_i, 0) &= 1 + \text{Ind}(s_i, P) && \text{in case (2);} \\
 \text{Ind}(f(s_i) - s_i, 0) &= \text{Ind}(s_i, P) && \text{in case (3).}
 \end{aligned}$$

Let N_1 be the number of repelling fixed points which were originally on ∂V , and N_2 be the number of arcs s_i of type (2). Observe that $N_1 = \sum n_i$ (where the sum is over the indices for which n_i is defined). Summing up, we have that

$$\begin{aligned}
 \text{Ind}(f(\Gamma_{\widehat{V}}) - \Gamma_{\widehat{V}}, 0) &= 1 + N_1 + N_2 + \sum_i \text{Ind}(s_i, P) \\
 &= 1 + N_1 + N_2 + \text{Ind}(\Gamma_{\widehat{V}}, P) \\
 &= 1 + N_1 + N_2,
 \end{aligned}$$

where we have used that P is contained in the unbounded connected component of $\mathbb{C} \setminus \Gamma_{\widehat{V}}$, so $\text{Ind}(\Gamma_{\widehat{V}}, P) = 0$. We deduce from this that there is exactly one interior fixed point. Indeed,

- there are exactly N_1 repelling fixed points which have been included in the region due to the modifications described in Lemma 6.1 (we do not need to count the parabolic ones, since they are no longer in V' due to Lemma 6.2);
- for each s_i of type (2) there is exactly one fixed point in the interior \widehat{V} which is the landing point of the unique fixed ray asymptotically contained in that fundamental domain (notice that those rays must land alone or otherwise they would be part of a ray pair).

This means that there are $N_1 + N_2$ fixed points in \widehat{V} which are landing points of fixed rays, hence exactly one fixed point is left. This fixed point must be interior because there are no more fixed rays left to land on it and because we assumed that V (and hence \widehat{V}) contains no virtual fixed points at all.

To conclude the proof of the Main Theorem we have to take into account the basic regions containing virtual fixed points. We will use the global counting in Theorem 5.2 and follow an argument analogous to the one in [GM, Proof of Theorem 3.3].

Let $N = \#\mathcal{F}$, and n be the number of landing points of fixed rays which are asymptotically contained in some $F \in \mathcal{F}$. By Theorem 5.2 there are $N + 1$ fixed points, counted with multiplicity, which are landing points of rays asymptotically contained in some $F \in \mathcal{F}$, or interior fixed points. Hence there are $N + 1 - n$ fixed points which therefore are interior or virtual.

On the other hand, let m be the total number of basic regions and v be the number of basic regions containing at least one virtual fixed point. It is easy to check that $m = N - n + 1$. Hence $m - v = N - n + 1 - v$ regions contain no virtual fixed point. Thus by the index counting above, there are $N - n + 1 - v$ interior fixed points, one inside each of these regions. Now let $v' \geq v$ be the actual number of virtual fixed points. Then the total number of fixed points which are interior or virtual is at least $N - n + 1 - v + v'$ (notice that *a priori* there could be interior fixed points in the v regions with at least one virtual fixed point).

Putting together both computations we obtain that

$$N + 1 - n \geq N - n + 1 - v + v'.$$

Hence $v' \leq v$ from which we deduce that $v' = v$. This implies that there is exactly one virtual fixed point in each of the v regions which had at least one of them. But moreover, all fixed points have been accounted for, so these v regions contain no interior fixed point.

7 Proof of corollaries D and E

As explained in the introduction, Theorem C is an immediate consequence of the Main Theorem, replacing f by f^p and observing that class $\widehat{\mathcal{B}}$ is closed under composition. In this section we prove some of the many corollaries of Theorem C. We recall that the period is not assumed to be exact.

Corollary 7.1. *If f is a function in $\widehat{\mathcal{B}}$ whose periodic rays land,*

1. *There are only finitely many interior periodic points of any given period. In particular, there are only finitely many non-repelling cycles of any given period.*
2. *Any two periodic Fatou components can be separated by a periodic ray pair.*
3. *There are no Cremer periodic points on the boundary of periodic Fatou components.*
4. *If z_0 is a parabolic point, for each repelling petal there is at least one ray which lands at z_0 through that repelling petal. In particular, any two of its attracting petals are separated by a ray pair of the same period.*
5. *For any given period p , there are only finitely many (possibly none) periodic points of period p which are landing points of more than one periodic ray. None of them is the landing point of infinitely many rays of the same period.*

Proof of Corollary 7.1. 1. By Theorem C, for each p there are finitely many basic regions for f^p , and each of them contains at most one interior fixed point of f^p which is an interior periodic point of period p (not necessarily exact) for f .

2. Let U_1, U_2 be two periodic Fatou components of least common period p . Then $\overline{U_1}$ and $\overline{U_2}$ each contain a point z_1, z_2 respectively which is fixed under f^p . If both z_1 and z_2 are in the interior of their Fatou component, they are interior periodic points, so they are separated by a p -periodic ray pair by Theorem C. As dynamic rays cannot intersect Fatou components, and Fatou components are connected sets, it follows that the same ray pair separates U_1 from U_2 . If instead at least one of them is on the boundary, say z_1 , then z_1 is a parabolic point and U_1 is a virtual fixed point for f^p . In particular, by Theorem C, there are no other interior p -periodic points or virtual fixed points of f^p in the basic region V containing U_1 , so U_2 is separated from U_1 by two rays on the boundary of V (observe that U_2 could also be another virtual fixed point of f^p attached to the same parabolic point if $z_1 = z_0$).
3. Observe that Cremer points are interior periodic points, since by the Snail Lemma (see Lemma 16.2 in [Mi]) no periodic rays can land at Cremer points. Hence they can be separated from any other interior periodic point or periodic Fatou component as in the previous case.
4. If z_0 is a multiple fixed point for f^p which has only one immediate basin attached, then there is at least a fixed ray of f^p landing at z_0 , otherwise z_0 would be an interior fixed point and would be contained in the same basic region as its attracting petal, contradicting Theorem C. If z_0 has more than one virtual fixed point of f^p attached, and there were a repelling petal which does not contain any fixed ray for f^p landing at z_0 , then its two adjacent immediate basins (virtual fixed points) would be contained in the same basic region for f^p , contradicting again Theorem C.
5. If not, there would be infinitely many basic regions for f^p , contradicting Proposition A. □

From the fact that Fatou components can be separated by periodic ray pairs we obtain the following additional corollary. Given an invariant Siegel disc Δ , we say that U is a hidden component of Δ if U is a bounded connected component of $\mathbb{C} \setminus \overline{\Delta}$. The proof of the next corollary follows the outline of Lemmas 2, 3 and 10 in [CR]; see also [Ro].

Corollary 7.2. *If $f \in \widehat{\mathcal{B}}$ and all periodic rays land, any hidden component of a bounded invariant Siegel disc is either a wandering domain or preperiodic to the Siegel disc itself.*

Proof. Observe that hidden components are bounded Fatou components. In fact, $\partial\Delta$ is bounded and forward invariant, and $\partial U \subset \partial\Delta$, so by the Maximum Principle and Montel's Theorem U is contained in the Fatou set. Suppose that there is a hidden component U which is not preperiodic to Δ . Observe that $\partial f(U) \subset f(\partial U)$ because f is holomorphic hence open, and $f(U) \cap \partial\Delta = \emptyset$ because $f(U) \subset F(f)$ and $\partial\Delta \subset J(f)$. Also $f(U)$ is bounded and connected because it is the image of a bounded connected set, and its boundary is contained in $\partial\Delta$, so $f(U)$ is also a hidden component of Δ . Hence, if U is not a wandering domain, it is preperiodic to some periodic Fatou component \tilde{U} which is a hidden component of Δ ; by part 4. of Corollary 7.1, unless $\tilde{U} = \Delta$, they can be separated by two periodic rays landing together, hence $\partial\tilde{U} \cap \partial\Delta$ is at most a single point- which is not possible as $\partial\tilde{U} \subset \partial\Delta$. \square

References

- [Ba] K. Baranski, *Trees and hairs for some hyperbolic entire maps of finite order*, Math. Z. **257** (2007), 33–59.
- [Bi] C. J. Bishop, *Constructiong entire functions by quasiconformal folding*. Preprint. <http://www.math.sunysb.edu/~bishop/papers/classS.pdf>
- [CR] A. Chéritat, P. Roesch, *Herman's condition and Siegel discs of polynomials*, Preprint (2011), ArXiv:111.4629v1.
- [DH] A. Douady, J. H. Hubbard, *Étude dynamique des polynômes complexes*, Prépublications mathématiques d'Orsay (1984/1985), no. 2/4.
- [DT] R. L. Devaney, F. Tangerman, *Dynamics of entire functions near the essential singularity*, Erg. Th. and Dyn. Syst. **6** (1986), no.4, 489–503.
- [EL] A. Eremenko, M. Lyubich, *Dynamical properties of some classes of entire functions*, Ann. Inst. Fourier (Grenoble) **42** (1992), no. 4, 989–1020.
- [Fa] N. Fagella, *Dynamics of the complex standard family*, J. Math. Anal. Appl. **229** (1999), no.1, 1–31.
- [GK] L. Goldberg, L. Keen, *A finiteness theorem for a dynamical class of entire functions*, Erg. Th. and Dyn. Syst. **6** (1986) no.2, 183–192.
- [GM] L. Goldberg, J. Milnor, *Fixed points of polynomial maps. Part II. Fixed point portraits*, Ann. Sci. École Norm. Sup. (4) **26** (1993), no. 1, 51–98.

- [Mi] J. Milnor, *Dynamics in one complex variable*, Annals of Mathematics Studies, Princeton University Press, 2006.
- [Rel] L. Rempe, *Siegel discs and periodic rays of entire functions*, J. Reine Angew. Math. **624** (2008), 81–102.
- [R³S] G. Rottenfusser, J. Rückert, L. Rempe, D. Schleicher, *Dynamic rays of bounded-type entire functions*, Ann. Math. **173** (2010), 77–125.
- [Ro] J. T. Rogers, *Diophantine conditions imply critical points on the boundaries of Siegel discs of polynomials*, Comm. Math. Phys. **195** (1998), 175–193.
- [Why] G. T. Whyburn, *Topological Analysis*, Princeton University Press, 1958.